

Lower-Dimensional Superstrings in the Double-Spinor Formalism

Hiroshi KUNITOMO¹ and Shun'ya MIZOGUCHI^{2,3}

¹*Yukawa Institute for Theoretical Physics, Kyoto University
 Kyoto 606-8502, Japan*

²*High Energy Accelerator Research Organization (KEK)
 Tsukuba 305-0801, Japan*

³*Department of Particle and Nuclear Physics
 The Graduate University for Advanced Studies
 Tsukuba 305-0801, Japan*

Abstract

We study lower-dimensional superstrings in the double-spinor formalism introduced by Aisaka and Kazama. These superstrings can be consistently quantized and are equivalent to the lower-dimensional pure-spinor superstrings proposed by Grassi and Wyllard. The unexpected physical spectrum of the pure-spinor superstrings may thus be regarded as a manifestation of noncriticality. We also discuss how to couple these covariant superstrings to the compactified degrees of freedom described by the $N = 2$ superconformal field theory.

§1. Introduction

The pure-spinor (PS) formalism, proposed by Berkovits, allows quantization of superstrings in a manner that preserves the manifest super-Poincaré covariance.¹⁾ Defined basically as a free conformal field theory (CFT), it provides a powerful framework for computing multipoint/multiloop amplitudes and for studying superstrings in RR backgrounds, which cannot be realized with other formalisms.

One of the unsolved problems of the PS formalism is to elucidate how the concept of “criticality” can be understood in this framework. In the RNS formalism, the critical dimension is that in which conformal anomalies cancel and the BRST charge becomes nilpotent, while in the light-cone Green-Schwarz (GS) formalism, this is the only dimension in which the global Lorentz invariance is unbroken. By contrast, the lower-dimensional versions of the PS formalism have been constructed^{2),3)} by simply anticipating analogous free CFTs with some plausible “pure spinor” conditions,^{*)} in which the BRST charge is exactly nilpotent and the Lorentz algebra has no anomaly. Therefore, they seem to be consistent theories, even quantum mechanically, although they also have some unexpected features such as the appearance of an *off-shell* vector multiplet in the open string spectrum at the lowest level. Like the original PS formalism, these theories are not based on Lagrangians, and this makes it difficult to understand their fundamental nature, and whether or not (and if so how) they are related to lower-dimensional (non-critical) superstrings in other formalisms.

Recently, a Lagrangian formulation of the ($D = 10$) PS superstring was proposed by Aisaka and Kazama.⁴⁾ A special feature of their formulation is that it involves, in addition to the ordinary superspace coordinates (X^μ, θ^α) in the GS superstring, another fermionic field, $\tilde{\theta}^\alpha$. The Lagrangian of this double-spinor (DS) formalism possesses world-sheet reparametrization invariance, a manifest space-time super Poincaré symmetry and a new local fermionic symmetry, which can be used to gauge away one of the fermionic fields. Since the Lagrangian of the DS formalism is reduced to the original GS Lagrangian by setting $\tilde{\theta}^\alpha$ to zero, this guarantees, at least classically, the equivalence of the DS superstring and the GS superstring.

The DS superstring can be quantized without difficulty using the conventional canonical BRST method. Surprisingly, the authors of 4) have shown that fields which have nontrivial Dirac brackets can be redefined in such a way that all of them become free fields. Therefore, the quantized theory can be described by a simple conformal field theory represented by the free fields. It is also proved in 4) that the physical spectrum of this DS superstring coincides with that of the PS superstring.

^{*)} It should be noted that this is in fact not the pure-spinor condition in lower dimensions.

In this paper, to gain a better understanding of the mysterious features of the lower-dimensional PS superstrings, we study the corresponding lower-dimensional (actually $d = 4$ and $d = 6$) DS superstrings. The equivalence to the lower-dimensional GS superstrings is manifest for the same reason as in the ten-dimensional case. We show that the lower-dimensional DS superstrings are also equivalent to the lower-dimensional PS superstrings. Therefore, it is suggested that the unexpected physical spectrum of the PS superstrings can be interpreted as a manifestation of “noncriticality” in the PS formalism. Then we search for a possible method for coupling these theories to the degrees of freedom of some compactified spaces, which we call the Calabi-Yau (CY) sector. It is well known that such degrees of freedom can be described by unitary representations of the $N = 2$ superconformal field theory with $c = 9$ (6) for $d = 4$ (6). We attempt to combine these two sectors in two different manners. Unfortunately, however, the spectrum in the resultant combined system is a tensor product of the physical spectrum of two sectors and thus is not an expected on-shell spectrum.

The paper is organized as follows. In §2 we briefly review the $d = 4$ and $d = 6$ PS superstrings^{2),3)} defined by naive extensions of the ten-dimensional PS superstring. It is shown that the BRST charges are nilpotent without anomaly, but the lowest-level physical state is an off-shell vector multiplet. The four-dimensional superstring in the DS formalism is studied in §3. It is possible to carry out the quantization in a manner parallel to that for the ten-dimensional case. The physical spectrum is shown to be equivalent to the PS formalism in four dimensions. A similar argument for the six-dimensional case is given in §4. The coupling to the degrees of freedom of the compactified space is discussed in §5. We attempt to apply two different methods for combining the two sectors, but the physical spectrum is simply given by tensor products of the unexpected, off-shell, spectrum of the lower-dimensional superstring with the (anti-)chiral ring of the CY sector. Finally, in §6, we summarize our results and discuss some remaining problems. The appendix contains a summary of our notation and spinor conventions.

§2. PS formalism in lower dimensions

As in $D = 10$, the lower-dimensional PS superstring is defined by free fields describing a map from the world-sheet to the target superspace. The bosonic coordinates $X^\mu(z)$ satisfy the free field operator product^{*)}

$$X^\mu(z)X^\nu(w) \sim \eta^{\mu\nu} \log(z - w), \quad (2.1)$$

^{*)} In this section, we concentrate only on the holomorphic sector, or open strings. It is easy to combine this sector with the anti-holomorphic sector to get closed strings.

where $\mu, \nu = 0, 1, \dots, d-1$. The fermionic coordinates are described by a first-order form of conformal dimension $(1, 0)$, whose explicit expression depends strongly on the space-time dimensionality. The spinor conventions and the notation used in this paper are summarized in the appendix A.

2.1. Four-dimensional PS superstring

As is well known, the fermionic coordinates of four-dimensional superspace consist of a pair of complex conjugate Weyl spinors, θ^α and $\bar{\theta}^{\dot{\alpha}}$. Introducing their conjugate fields, p_α and $\bar{p}_{\dot{\alpha}}$, we assume the free field operator products

$$\theta^\alpha(z)p_\beta(w) \sim \frac{\delta^\alpha_\beta}{z-w}, \quad \bar{\theta}^{\dot{\alpha}}(z)\bar{p}_{\dot{\beta}}(w) \sim \frac{\delta^{\dot{\alpha}}_{\dot{\beta}}}{z-w}. \quad (2.2)$$

We assign θ^α and $\bar{\theta}^{\dot{\alpha}}$ conformal dimension 0 and p_α and $\bar{p}_{\dot{\alpha}}$ conformal dimension 1. In addition to these fields, we also introduce the bosonic spinor fields λ^α and $\bar{\lambda}^{\dot{\alpha}}$ satisfying the *pure-spinor constraint*

$$\lambda\sigma^\mu\bar{\lambda} = 0. \quad (2.3)$$

Using these pure-spinor fields, the BRST charge is defined by

$$Q_{PS} = \oint \frac{dz}{2\pi i} (\lambda^\alpha d_\alpha + \bar{\lambda}^{\dot{\alpha}} \bar{d}_{\dot{\alpha}}). \quad (2.4)$$

Here d_α and $\bar{d}_{\dot{\alpha}}$ are the currents corresponding to the super-covariant derivatives

$$d_\alpha = p_\alpha - i\partial X^\mu (\sigma_\mu \bar{\theta})_\alpha - \frac{1}{2} ((\theta\sigma^\mu\partial\bar{\theta}) - (\partial\theta\sigma^\mu\bar{\theta})) (\sigma_\mu \bar{\theta})_\alpha, \quad (2.5a)$$

$$\bar{d}_{\dot{\alpha}} = \bar{p}_{\dot{\alpha}} - i\partial X^\mu (\theta\sigma_\mu)_{\dot{\alpha}} - \frac{1}{2} ((\theta\sigma^\mu\partial\bar{\theta}) - (\partial\theta\sigma^\mu\bar{\theta})) (\theta\sigma_\mu)_{\dot{\alpha}}, \quad (2.5b)$$

which satisfy

$$d_\alpha(z)\bar{d}_{\dot{\alpha}}(w) \sim \frac{2i(\sigma_\mu)_{\alpha\dot{\alpha}}\pi^\mu(w)}{z-w}, \quad (2.6)$$

with

$$\pi^\mu = i\partial X^\mu + (\theta\sigma^\mu\partial\bar{\theta}) - (\partial\theta\sigma^\mu\bar{\theta}). \quad (2.7)$$

The BRST charge Q_{PS} is nilpotent, due to the pure-spinor constraint (2.3), even at the quantum level. The physical states are defined as the cohomology of this BRST charge. However, it possesses a rather unexpected spectrum, because the on-shell condition is not imposed. For example, the lowest-level vertex operator with ghost number 1, which is expected to involve the physical state, has the form

$$W = \lambda^\alpha A_\alpha(x, \theta, \bar{\theta}) + \bar{\lambda}^{\dot{\alpha}} A_{\dot{\alpha}}(x, \theta, \bar{\theta}), \quad (2.8)$$

where A_α and $A_{\dot{\alpha}}$ are conventional superfields of zero-modes $(x^\mu, \theta^\alpha, \bar{\theta}^{\dot{\alpha}})$. Then the BRST invariance $\{Q, W\} = 0$ implies the conditions

$$D_{(\alpha} A_{\beta)}(x, \theta, \bar{\theta}) = 0, \quad (2.9a)$$

$$\bar{D}_{(\dot{\alpha}} A_{\dot{\beta}})(x, \theta, \bar{\theta}) = 0, \quad (2.9b)$$

$$D_\alpha A_{\dot{\alpha}}(x, \theta, \bar{\theta}) + \bar{D}_{\dot{\alpha}} A_\alpha(x, \theta, \bar{\theta}) + 2i(\sigma^\mu)_{\alpha\dot{\alpha}} A_\mu(x, \theta, \bar{\theta}) = 0, \quad (2.9c)$$

with an arbitrary superfield A_μ . These are the well-known torsion constraints⁵⁾ and can be solved in terms of a real superfield $V(x, \theta, \bar{\theta})$ as

$$A_\alpha = iD_\alpha V, \quad A_{\dot{\alpha}} = -i\bar{D}_{\dot{\alpha}} V, \quad A_\mu = \frac{1}{4}(\bar{\sigma}_\mu)^{\dot{\alpha}\alpha}[D_\alpha, \bar{D}_{\dot{\alpha}}]V, \quad (2.10)$$

up to BRST exact pieces. There are no further restrictions, and thus the lowest-level physical states constitute an *off-shell* vector multiplet.

2.2. Six-dimensional PS superstring

The fermionic coordinates of six-dimensional superspace are $SU(2)$ -Majorana-Weyl (MW) spinors θ_I^α ($I = 1, 2$). The free field operator product with the conjugate field p_α^I is

$$\theta_I^\alpha(z)p_\beta^J(w) \sim \frac{\delta_\beta^\alpha \delta_I^J}{z-w}. \quad (2.11)$$

We assign θ_I^α and p_α^I conformal dimension 0 and 1, respectively. The bosonic pure-spinors λ_α^I satisfy

$$\lambda_I C \gamma^\mu \lambda^I = 0. \quad (2.12)$$

Then the BRST charge is defined by

$$Q_{PS} = \oint \frac{dz}{2\pi i} \lambda_I^\alpha d_\alpha^I, \quad (2.13)$$

which is also nilpotent quantum mechanically, since the operator product relations

$$d_\alpha^I(z)d_\beta^J(w) \sim \frac{2i\epsilon^{IJ}(C\gamma_\mu)_{\alpha\beta}\pi^\mu(w)}{z-w} \quad (2.14)$$

hold between the super-covariant currents

$$d_\alpha^I = p_\alpha^I + i\partial X^\mu (C\gamma_\mu \theta^I)_\alpha + \frac{1}{2}(\theta^K C \gamma^\mu \partial \theta_K)(C\gamma_\mu \theta^I)_\alpha, \quad (2.15a)$$

$$\pi^\mu = i\partial X^\mu + (\theta^K C \gamma^\mu \partial \theta_K). \quad (2.15b)$$

The lowest-level vertex operator with ghost number 1 in six dimensions is now given by

$$W = \lambda_I^\alpha A_\alpha^I(x, \theta_I), \quad (2.16)$$

where A_α is a superfield of the zero-modes (x^μ, θ_I^α) . The BRST invariance $\{Q, W\} = 0$ yields the conditions

$$D_{(\alpha}^I A_{\beta)}^J = 0, \quad (2.17a)$$

$$D_{I[\alpha} A_{\beta]}^I + 2i(C\gamma^\mu)_{\alpha\beta} A_\mu = 0, \quad (2.17b)$$

with an arbitrary superfield A_μ . These six-dimensional torsion constraints can also be solved similarly to those in the four-dimensional case.^{2),6)} The lowest-level physical states are therefore also given by a six-dimensional *off-shell* vector multiplet.

§3. DS formalism in four dimensions

The Lagrangian formulation of the PS superstring is given in 4). This formulation can be easily extended to the lower-dimensional case.

3.1. Lagrangian, symmetries and constraints

The four-dimensional superstring in the DS formalism is defined by using the super-space coordinates $(x^\mu, \theta^{A\alpha}, \bar{\theta}^{A\dot{\alpha}})$ and the additional fermionic fields $(\tilde{\theta}^{A\alpha}, \tilde{\bar{\theta}}^{A\dot{\alpha}})$ ($A=1,2$). The Lagrangian is then given by

$$\mathcal{L} = \mathcal{L}_K + \mathcal{L}_{WZ}, \quad (3.1a)$$

$$\mathcal{L}_K = -\frac{1}{2}\sqrt{-g}g^{ab}\Pi_a^\mu\Pi_{\mu b}, \quad (3.1b)$$

$$\mathcal{L}_{WZ} = \epsilon^{ab}\Pi_a^\mu(W_{\mu b} - \hat{W}_{\mu b}) - \epsilon^{ab}W_a^\mu\hat{W}_{\mu b}, \quad (3.1c)$$

where

$$\Pi_a^\mu = \partial_a X^\mu - \sum_{A=1}^2 i\partial_a(\theta^A\sigma^\mu\tilde{\bar{\theta}}^A - \tilde{\theta}^A\sigma^\mu\bar{\theta}^A) - \sum_{A=1}^2 W_a^{A\mu}, \quad (3.2)$$

$$W_a^{A\mu} = i\Theta^A\sigma^\mu\partial_a\bar{\Theta}^A - i\partial_a\Theta^A\sigma^\mu\bar{\Theta}^A, \quad (3.3)$$

$$\Theta^A = \tilde{\theta}^A - \theta^A, \quad \bar{\Theta}^A = \tilde{\bar{\theta}}^A - \bar{\theta}^A. \quad (3.4)$$

We use the same notation as in Ref. 4); for example, $W_a^{1\mu} = W_a^\mu$, $W_a^{2\mu} = \hat{W}_a^\mu$, etc. This Lagrangian is invariant under the world-sheet reparametrization and the space-time Poincaré transformation, where the space-time supersymmetry transformation is defined by

$$\delta\theta^A = \epsilon^A, \quad \delta\bar{\theta}^A = \bar{\epsilon}^A, \quad (3.5a)$$

$$\delta\tilde{\theta}^A = 0, \quad \delta\tilde{\bar{\theta}}^A = 0, \quad (3.5b)$$

$$\delta X^\mu = \sum_{A=1}^2 i(\epsilon^A\sigma^\mu\bar{\theta}^A - \theta^A\sigma^\mu\bar{\epsilon}^A). \quad (3.5c)$$

The additional fermionic fields are inert under this supersymmetry.

The Lagrangian has another important symmetry, the local supersymmetry, given by

$$\delta\theta^A = \chi^A, \quad \delta\tilde{\theta}^A = \chi^A, \quad (3.6a)$$

$$\delta\bar{\theta}^A = \bar{\chi}^A, \quad \delta\tilde{\bar{\theta}}^A = \bar{\chi}^A, \quad (3.6b)$$

$$\delta X^\mu = \sum_{A=1}^2 i(\chi^A \sigma^\mu \bar{\Theta}^A - \Theta^A \sigma^\mu \bar{\chi}^A), \quad (3.6c)$$

which guarantees the equivalence to the conventional Green-Schwarz formalism. Using this local symmetry, we can set the additional fermionic fields as $\tilde{\theta}^{A\alpha} = \tilde{\bar{\theta}}^{A\dot{\alpha}} = 0$. Then, the Lagrangian (3.1) becomes that of the Green-Schwarz formalism.⁷⁾

Let us consider the canonical quantization of the Lagrangian (3.1). First, the canonical conjugate of X^μ can be computed as

$$k_\mu = -\sqrt{-g}g^{0b}\Pi_{\mu b} + W_{\mu 1} - \hat{W}_{\mu 1}. \quad (3.7)$$

Then, computing the canonical conjugates $k_\alpha^A, \bar{k}_{\dot{\alpha}}^A, \tilde{k}_\alpha^A$ and $\tilde{\bar{k}}_{\dot{\alpha}}^A$ of the fermionic fields $\theta^{A\alpha}, \bar{\theta}^{A\dot{\alpha}}, \tilde{\theta}^{A\alpha}$ and $\tilde{\bar{\theta}}^{A\dot{\alpha}}$, we obtain the primary constraints

$$D_\alpha^A = k_\alpha^A + i(\not{k}\tilde{\bar{\theta}}^A)_\alpha + i(k^\mu + \eta_A(\Pi_1^\mu + W_1^{\mu\bar{A}}))(\sigma_\mu \bar{\Theta}^A)_\alpha \approx 0, \quad (3.8a)$$

$$\bar{D}_\alpha^A = \bar{k}_{\dot{\alpha}}^A + i(\tilde{\bar{\theta}}^A \not{k})_{\dot{\alpha}} + i(k^\mu + \eta_A(\Pi_1^\mu + W_1^{\mu\bar{A}}))(\Theta^A \sigma_\mu)_{\dot{\alpha}} \approx 0, \quad (3.8b)$$

$$\tilde{D}_\alpha^A = \tilde{k}_\alpha^A - i(\not{k}\tilde{\bar{\theta}}^A)_\alpha - i(k^\mu + \eta_A(\Pi_1^\mu + W_1^{\mu\bar{A}}))(\sigma_\mu \bar{\Theta}^A)_\alpha \approx 0, \quad (3.8c)$$

$$\tilde{\bar{D}}_\alpha^A = \tilde{\bar{k}}_{\dot{\alpha}}^A - i(\tilde{\theta}^A \not{k})_{\dot{\alpha}} - i(k^\mu + \eta_A(\Pi_1^\mu + W_1^{\mu\bar{A}}))(\Theta^A \sigma_\mu)_{\dot{\alpha}} \approx 0, \quad (3.8d)$$

where $\eta_1 = -\eta_2 = 1$ and $\bar{A} = 2(1)$ for $A = 1(2)$. For later use, it is convenient to define $\Delta_\alpha = D_\alpha + \tilde{D}_\alpha$, $\bar{\Delta}_{\dot{\alpha}} = \bar{D}_{\dot{\alpha}} + \tilde{\bar{D}}_{\dot{\alpha}}$.

Using the ADM decomposition

$$g_{ab} = \begin{pmatrix} -N^2 + \gamma(N^1)^2 & \gamma N^1 \\ \gamma N^1 & \gamma \end{pmatrix}, \quad (3.9)$$

we obtain the Hamiltonian as

$$\begin{aligned} \mathcal{H} &= \frac{N}{\sqrt{\gamma}} \frac{1}{2} \left((k_\mu - W_{\mu 1} + \hat{W}_{\mu 1})(k^\mu - W_1^\mu + \hat{W}_1^\mu) + \Pi_1^\mu \Pi_{\mu 1} \right) \\ &\quad + N^1 (k_\mu - W_{\mu 1} + \hat{W}_{\mu 1}) \Pi_1^\mu + \dot{\theta}^{A\alpha} D_\alpha^A + \dot{\bar{\theta}}^{A\dot{\alpha}} \bar{D}_{\dot{\alpha}}^A + \dot{\tilde{\theta}}^{A\alpha} \tilde{D}_\alpha^A + \dot{\tilde{\bar{\theta}}}^{A\dot{\alpha}} \tilde{\bar{D}}_{\dot{\alpha}}^A, \\ &\equiv \frac{N}{\sqrt{\gamma}} T_0 + N^1 T_1 + \dot{\theta}^{A\alpha} D_\alpha^A + \dot{\bar{\theta}}^{A\dot{\alpha}} \bar{D}_{\dot{\alpha}}^A + \dot{\tilde{\theta}}^{A\alpha} \tilde{D}_\alpha^A + \dot{\tilde{\bar{\theta}}}^{A\dot{\alpha}} \tilde{\bar{D}}_{\dot{\alpha}}^A, \end{aligned} \quad (3.10)$$

where the energy-momentum tensors are given by

$$T_+ = \frac{1}{2}(T_0 + T_1) = \frac{1}{4}\Pi^\mu \Pi_\mu, \quad (3.11a)$$

$$T_- = \frac{1}{2}(T_0 - T_1) = \frac{1}{4}\hat{\Pi}^\mu \hat{\Pi}_\mu, \quad (3.11b)$$

with

$$\begin{aligned} \Pi^\mu &= k^\mu - W_1^\mu + \hat{W}_1^\mu + \Pi_1^\mu, \\ &= k^\mu + X'^\mu - \sum_A i(\theta^A \sigma^\mu \tilde{\theta}^A - \tilde{\theta}^A \sigma^\mu \bar{\theta}^A)' - 2W_1^\mu, \end{aligned} \quad (3.12a)$$

$$\begin{aligned} \hat{\Pi}^\mu &= k^\mu - W_1^\mu + \hat{W}_1^\mu - \Pi_1^\mu, \\ &= k^\mu - X'^\mu + \sum_A i(\theta^A \sigma^\mu \tilde{\theta}^A - \tilde{\theta}^A \sigma^\mu \bar{\theta}^A)' + 2\hat{W}_1^\mu. \end{aligned} \quad (3.12b)$$

Next we set the canonical Poisson brackets as

$$\{X^\mu(\sigma), k^\nu(\sigma')\}_P = \eta^{\mu\nu} \delta(\sigma - \sigma'), \quad (3.13a)$$

$$\{\theta^{A\alpha}(\sigma), k_\beta^B(\sigma')\}_P = -\delta^{AB} \delta_\beta^\alpha \delta(\sigma - \sigma'), \quad (3.13b)$$

$$\{\bar{\theta}^{A\dot{\alpha}}(\sigma), \bar{k}_\beta^B(\sigma')\}_P = -\delta^{AB} \delta_\beta^{\dot{\alpha}} \delta(\sigma - \sigma'), \quad (3.13c)$$

$$\{\tilde{\theta}^{A\alpha}(\sigma), \tilde{k}_\beta^B(\sigma')\}_P = -\delta^{AB} \delta_\beta^\alpha \delta(\sigma - \sigma'), \quad (3.13d)$$

$$\{\tilde{\bar{\theta}}^{A\dot{\alpha}}(\sigma), \tilde{\bar{k}}_\beta^B(\sigma')\}_P = -\delta^{AB} \delta_\beta^{\dot{\alpha}} \delta(\sigma - \sigma'). \quad (3.13e)$$

As in the ten-dimensional case,⁴⁾ the algebra of constraints can be separated into left ($A = 1$, or un-hatted) and right ($A = 2$, or hatted) sectors, although the generators include both un-hatted and hatted fields. Thus, for simplicity, we hereafter concentrate on the left sector. Using the canonical Poisson brackets (3.13), we can compute the Poisson brackets among the fermionic constraint generators ($\Delta_\alpha, \bar{\Delta}_{\dot{\alpha}}, \tilde{D}_\alpha, \tilde{\bar{D}}_{\dot{\alpha}}$) as

$$\{\tilde{D}_\alpha(\sigma), \tilde{\bar{D}}_{\dot{\alpha}}(\sigma')\}_P = 2i\Pi^\mu(\sigma)(\sigma_\mu)_{\alpha\dot{\alpha}}\delta(\sigma - \sigma'), \quad (3.14)$$

with all others vanishing.

In addition to these fermionic constraints, we also obtain the Virasoro constraint, $T_+ \approx 0$, for the left sector, from the Hamiltonian (3.10) as the secondary constraint. It is convenient to define the total energy-momentum tensor

$$T = \frac{1}{4}\Pi^\mu \Pi_\mu + \Theta'^\alpha \tilde{D}_\alpha + \bar{\Theta}'^{\dot{\alpha}} \tilde{\bar{D}}_{\dot{\alpha}} + \theta^\alpha \Delta_\alpha + \bar{\theta}^{\dot{\alpha}} \bar{\Delta}_{\dot{\alpha}}, \quad (3.15)$$

by adding the fermion contributions, which are also weakly vanishing, due to the constraint (3.8). The Poisson bracket of this tensor is that of the Virasoro algebra:

$$\{T(\sigma), T(\sigma')\}_P = 2T(\sigma)\delta'(\sigma - \sigma') + T'(\sigma)\delta(\sigma - \sigma'). \quad (3.16)$$

Now, let us study the constraint algebra (3.14). Due to the Virasoro constraints $T_+ \approx 0$, this implies that half of the constraints $\tilde{D}_\alpha \approx 0$ and $\tilde{\tilde{D}}_\alpha \approx 0$ are first class and the half are second class. As is well known, these constraints cannot be separated covariantly, and therefore we use the light-cone decomposition. The Poisson bracket

$$\{\tilde{D}_1(\sigma), \tilde{\tilde{D}}_1(\sigma')\}_P = 2i\Pi^+(\sigma)\delta(\sigma - \sigma') \quad (3.17)$$

shows that the first components $\tilde{D}_1$ and $\tilde{\tilde{D}}_1$ generate the second-class constraints. The other half of the constraints come from the κ -symmetry generated by

$$K = \tilde{D}_2 - \frac{\Pi}{\Pi^+} \tilde{D}_1, \quad (3.18a)$$

$$\bar{K} = \tilde{\tilde{D}}_2 - \frac{\bar{\Pi}}{\Pi^+} \tilde{\tilde{D}}_1, \quad (3.18b)$$

where $\Pi = \Pi^1 + i\Pi^2$ and $\bar{\Pi} = \Pi^1 - i\Pi^2$. These are first class, because they satisfy the relations

$$\{K(\sigma), \bar{K}(\sigma')\}_P = 8i(\mathcal{T} + \mathcal{K})(\sigma)\delta'(\sigma - \sigma') + (\tilde{D}_1, \tilde{\tilde{D}}_1)\text{-terms}, \quad (3.19)$$

where

$$\mathcal{T} = \frac{T}{\Pi^+}, \quad (3.20a)$$

$$\mathcal{K} = -\frac{1}{\Pi^+}(\Theta'^2 K + \bar{\Theta}'^2 \bar{K}). \quad (3.20b)$$

These relations are closed up to $(\tilde{D}_1, \tilde{\tilde{D}}_1)$ -terms, which are proportional to the second-class constraints $\tilde{D}_1 \approx 0$ and $\tilde{\tilde{D}}_1 \approx 0$ and can be set to zero after calculating the Dirac bracket. This is also consistent with the fact that the second-class constraints are transformed into themselves under this symmetry generated by the first-class constraints:

$$\{K(\sigma), \tilde{D}_1(\sigma')\}_P = 8i\frac{1}{\Pi^+}\bar{\Theta}'^2 \tilde{D}_1(\sigma)\delta(\sigma - \sigma'), \quad (3.21a)$$

$$\{K(\sigma), \tilde{\tilde{D}}_1(\sigma')\}_P = 0, \quad (3.21b)$$

$$\{\bar{K}(\sigma), \tilde{D}_1(\sigma')\}_P = 0, \quad (3.21c)$$

$$\{\bar{K}(\sigma), \tilde{\tilde{D}}_1(\sigma')\}_P = 8i\frac{1}{\Pi^+}\Theta'^2 \tilde{\tilde{D}}_1(\sigma)\delta(\sigma - \sigma'). \quad (3.21d)$$

The above decomposition of the constraints explicitly breaks the manifest Lorentz invariance. This breaking is, however, restricted to the sector of the additional fermionic fields, and therefore the space-time supersymmetry defined by (3.5) can be linearly realized.

If we choose the semi-light-cone gauge, defined by

$$\tilde{\theta}^2 \approx \tilde{\tilde{\theta}}^2 \approx 0, \quad (3.22)$$

to fix the κ -symmetry, all the constraints $\phi^I \approx 0$ with $\phi^I = (\tilde{D}_1, \tilde{\bar{D}}_1, K, \bar{K}, \tilde{\theta}^2, \tilde{\bar{\theta}}^2)$ become second class. We can take account of these second-class constraints by computing the Dirac bracket

$$\begin{aligned}
\{A(\sigma), B(\sigma')\}_D = & \{A(\sigma), B(\sigma')\}_P \\
& - \int d\sigma_1 \{A(\sigma), \tilde{D}_1(\sigma_1)\}_P \frac{1}{2i\Pi^+}(\sigma_1) \{\tilde{\bar{D}}_1(\sigma_1), B(\sigma')\}_P \\
& - \int d\sigma_1 \{A(\sigma), \tilde{\bar{D}}_1(\sigma_1)\}_P \frac{1}{2i\Pi^+}(\sigma_1) \{\tilde{D}_1(\sigma_1), B(\sigma')\}_P \\
& + 8i \int d\sigma_1 \{A(\sigma), \tilde{\theta}^2(\sigma_1)\}_P \mathcal{T}(\sigma_1) \{\tilde{\bar{\theta}}^2(\sigma_1), B(\sigma')\}_P \\
& + 8i \int d\sigma_1 \{A(\sigma), \tilde{\bar{\theta}}^2(\sigma_1)\}_P \mathcal{T}(\sigma_1) \{\tilde{\theta}^2(\sigma_1), B(\sigma')\}_P \\
& + \int d\sigma_1 \{A(\sigma), \tilde{\theta}^2(\sigma_1)\}_P \{K(\sigma_1), B(\sigma')\}_P \\
& + \int d\sigma_1 \{A(\sigma), K(\sigma_1)\}_P \{\tilde{\theta}^2(\sigma_1), B(\sigma')\}_P \\
& + \int d\sigma_1 \{A(\sigma), \tilde{\bar{\theta}}^2(\sigma_1)\}_P \{\bar{K}(\sigma_1), B(\sigma')\}_P \\
& + \int d\sigma_1 \{A(\sigma), \bar{K}(\sigma_1)\}_P \{\tilde{\bar{\theta}}^2(\sigma_1), B(\sigma')\}_P.
\end{aligned} \tag{3.23}$$

In this semi-light-cone gauge, the independent fields are $(\tilde{\theta}^1, \tilde{\bar{\theta}}^1, X^\mu, k_\mu, \theta^\alpha, k_\alpha, \bar{\theta}^{\dot{\alpha}}, \bar{k}_{\dot{\alpha}})$, which satisfy the Dirac brackets

$$\{\tilde{\theta}^1(\sigma), \tilde{\bar{\theta}}^1(\sigma')\}_D = \frac{i}{2\Pi^+}(\sigma) \delta(\sigma - \sigma'), \tag{3.24a}$$

$$\{X^\mu(\sigma), \tilde{\theta}^1(\sigma')\}_D = -\frac{1}{2\Pi^+}(\tilde{\theta}\sigma^\mu)_1(\sigma) \delta(\sigma - \sigma'), \tag{3.24b}$$

$$\{X^\mu(\sigma), \tilde{\bar{\theta}}^1(\sigma')\}_D = -\frac{1}{2\Pi^+}(\sigma^\mu \tilde{\bar{\theta}})_1(\sigma) \delta(\sigma - \sigma'), \tag{3.24c}$$

$$\{X^\mu(\sigma), X^\nu(\sigma')\}_D = \frac{i}{2\Pi^+} \left((\tilde{\theta}\sigma^\mu)_1(\sigma^\nu \tilde{\bar{\theta}})_1 - (\tilde{\theta}\sigma^\nu)_1(\sigma^\mu \tilde{\bar{\theta}})_1 \right) (\sigma) \delta(\sigma - \sigma'), \tag{3.24d}$$

$$\begin{aligned}
\{X^\mu(\sigma), k^\nu(\sigma')\}_D = & \eta^{\mu\nu} \delta(\sigma - \sigma') \\
& - \frac{i}{2\Pi^+} \left((\tilde{\theta}\sigma^\mu)_1(\sigma^\nu \bar{\Theta})_1 - (\Theta\sigma^\nu)_1(\sigma^\mu \tilde{\bar{\theta}}_1) \right) (\sigma) \delta'(\sigma - \sigma'),
\end{aligned} \tag{3.24e}$$

$$\{P^\mu(\sigma), k^\nu(\sigma')\}_D = -\frac{i}{2} \partial_\sigma \left(\frac{1}{\Pi^+} \left((\Theta\sigma^\mu)_1(\sigma^\nu \bar{\Theta})_1 - (\Theta\sigma^\nu)_1(\sigma^\mu \bar{\Theta})_1 \right) (\sigma) \delta'(\sigma - \sigma') \right), \tag{3.24f}$$

$$\{\tilde{\theta}^1(\sigma), k^\nu(\sigma')\}_D = -\frac{1}{2\Pi^+}(\Theta\sigma^\nu)_1(\sigma) \delta'(\sigma - \sigma'), \tag{3.24g}$$

$$\{\tilde{\bar{\theta}}^1(\sigma), k^\nu(\sigma')\}_D = -\frac{1}{2\Pi^+}(\sigma^\nu \bar{\Theta})_1(\sigma) \delta'(\sigma - \sigma'). \tag{3.24h}$$

These can be rewritten in simpler forms in the constraint plane defined by $\tilde{\theta}^2 = \tilde{\bar{\theta}}^2 = 0$. The non-trivial Dirac brackets are, for example,

$$\{\tilde{\theta}^1(\sigma), \tilde{\bar{\theta}}^1(\sigma')\}_D = \frac{i}{2\Pi^+}(\sigma)\delta(\sigma - \sigma'), \quad (3.25a)$$

$$\{X^-(\sigma), \tilde{\theta}^1(\sigma')\}_D = -\frac{1}{\Pi^+}\tilde{\theta}^1(\sigma)\delta(\sigma - \sigma'), \quad (3.25b)$$

$$\{X^-(\sigma), \tilde{\bar{\theta}}^1(\sigma')\}_D = -\frac{1}{\Pi^+}\tilde{\bar{\theta}}^1(\sigma)\delta(\sigma - \sigma'). \quad (3.25c)$$

We have now separated, as in 4), the first-class constraints from the second, the latter of which can be set to zero consistently with use of the Dirac bracket. With their non-trivial forms, it may appear difficult to quantize the theory. The authors of 4) found in $D = 10$ that one can redefine the independent fields so that the new ones satisfy simple free-field brackets. Here we show that we can do the same thing in the $D = 4$ DS theory.

First, we define

$$S = \sqrt{2\Pi^+}\tilde{\theta}^1, \quad \bar{S} = \sqrt{2\Pi^+}\tilde{\bar{\theta}}^1. \quad (3.26)$$

Then S and $\bar{S}$ become independent free fermions:

$$\{S(\sigma), \bar{S}(\sigma')\}_{D^*} = i\delta(\sigma - \sigma'), \quad (3.27a)$$

$$\{X^-(\sigma), S(\sigma')\}_{D^*} = 0, \quad (3.27b)$$

$$\{X^-(\sigma), \bar{S}(\sigma')\}_{D^*} = 0. \quad (3.27c)$$

Similarly for the remaining fields, we can find field redefinitions that make all of them free fields:

$$P^\mu = k^\mu - i(\tilde{\theta}\sigma^\mu\bar{\theta})' + i(\theta\sigma^\mu\tilde{\bar{\theta}})' + i(\hat{\theta}\sigma^\mu\hat{\bar{\theta}})' - i(\hat{\theta}\sigma^\mu\hat{\bar{\theta}})', \quad (3.28a)$$

$$p_1^A = k_1^A + \eta^A \left(-iX'^+\tilde{\bar{\theta}}^{A1} - 6(\theta^{A2}\tilde{\theta}'^{A2})\tilde{\bar{\theta}}^{A1} - 2(\theta^{A2}\bar{\theta}^{A2})\tilde{\bar{\theta}}'^{A1} \right), \quad (3.28b)$$

$$p_2^A = k_2^A + \eta^A \left(-iX'\tilde{\bar{\theta}}^{A1} - 6(\bar{\theta}'^{A2}\theta^{A1})\tilde{\bar{\theta}}^{A1} + 2(\bar{\theta}^{A2}\tilde{\theta}^{A1})'\tilde{\bar{\theta}}^{A1} \right. \\ \left. + 3(\bar{\theta}^A\bar{\theta}^A)'\tilde{\bar{\theta}}^{A1} + 2\bar{\theta}^{A2}(\tilde{\bar{\theta}}^{A1}\tilde{\theta}^{A1})' - 2(\bar{\theta}^{A2}\theta^{A1})\tilde{\bar{\theta}}'^{A1} + 2(\bar{\theta}^A\bar{\theta}^A)\tilde{\bar{\theta}}'^{A1} \right), \quad (3.28c)$$

$$\bar{p}_1^A = \bar{k}_1^A + \eta^A \left(-iX'^+\tilde{\bar{\theta}}^{A1} - 6(\bar{\theta}^{A2}\theta'^{A2})\tilde{\bar{\theta}}^{A1} - 2(\bar{\theta}^{A2}\theta^{A2})\tilde{\bar{\theta}}'^{A1} \right), \quad (3.28d)$$

$$\bar{p}_2^A = \bar{k}_2^A + \eta^A \left(-i\bar{X}'\tilde{\bar{\theta}}^{A1} - 6(\theta'^{A2}\bar{\theta}^{A1})\tilde{\bar{\theta}}^{A1} + 2(\theta^{A2}\tilde{\bar{\theta}}^{A1})'\tilde{\bar{\theta}}^{A1} \right. \\ \left. + 3(\theta^A\theta^A)'\tilde{\bar{\theta}}^{A1} + 2\theta^{A2}(\tilde{\bar{\theta}}^{A1}\tilde{\bar{\theta}}^{A1})' - 2(\theta^{A2}\bar{\theta}^{A1})\tilde{\bar{\theta}}'^{A1} + 2(\theta^A\theta^A)\tilde{\bar{\theta}}'^{A1} \right), \quad (3.28e)$$

which satisfy

$$\{X^\mu(\sigma), P^\nu(\sigma')\}_D = \eta^{\mu\nu}\delta(\sigma - \sigma'), \quad (3.29a)$$

$$\{\theta^{A\alpha}(\sigma), p_\beta^B(\sigma')\}_D = -\delta^{AB}\delta_\beta^\alpha\delta(\sigma - \sigma'), \quad (3.29b)$$

$$\{\bar{\theta}^{A\dot{\alpha}}(\sigma), \bar{p}_\beta^B(\sigma')\}_D = -\delta^{AB}\delta_\beta^\alpha\delta(\sigma - \sigma'). \quad (3.29c)$$

with the remaining brackets vanishing. These redefinitions also yield a complete separation⁴⁾ of the left and right sectors for the fundamental quantities, including the constraint generators. For instance, we have

$$\begin{aligned} \Pi^\mu &= P^\mu + X^{\mu'} \\ &\quad - 2i(\theta\sigma^\mu\bar{\theta}') + 2i(\theta'\sigma^\mu\bar{\theta}) - 2i(\tilde{\theta}\sigma^\mu\tilde{\theta}') + 2i(\tilde{\theta}'\sigma^\mu\tilde{\tilde{\theta}}) + 4i(\tilde{\theta}\sigma^\mu\bar{\theta}') - 4i(\theta'\sigma^\mu\tilde{\tilde{\theta}}), \end{aligned} \quad (3.30a)$$

$$\begin{aligned} \hat{\Pi}^\mu &= P^\mu - X^{\mu'} \\ &\quad + 2i(\hat{\theta}\sigma^\mu\hat{\theta}') - 2i(\hat{\theta}'\sigma^\mu\hat{\theta}) + 2i(\hat{\tilde{\theta}}\sigma^\mu\hat{\tilde{\theta}}') - 2i(\hat{\tilde{\theta}}'\sigma^\mu\hat{\tilde{\tilde{\theta}}}) - 4i(\hat{\tilde{\theta}}\sigma^\mu\hat{\theta}') + 4i(\hat{\theta}'\sigma^\mu\hat{\tilde{\tilde{\theta}}}). \end{aligned} \quad (3.30b)$$

Now let us rewrite the constraint generators in terms of these free fields. They become

$$D_1 = d_1 + i\sqrt{2\Pi^+}\bar{S}, \quad (3.31a)$$

$$D_2 = d_2 + i\sqrt{\frac{2}{\Pi^+}}\Pi\bar{S} + \frac{4}{\Pi^+}S\bar{S}(\bar{\theta}^2)', \quad (3.31b)$$

$$\bar{D}_1 = \bar{d}_1 + i\sqrt{2\Pi^+}S, \quad (3.31c)$$

$$\bar{D}_2 = \bar{d}_2 + i\sqrt{\frac{2}{\Pi^+}}\bar{\Pi}S - \frac{4}{\Pi^+}S\bar{S}(\theta^2)', \quad (3.31d)$$

$$\mathcal{T} = -\frac{1}{4}\frac{\Pi^\mu\Pi_\mu}{\Pi^+}, \quad (3.31e)$$

where d_α and $\bar{d}_{\dot{\alpha}}$ are the supercovariant spinor conjugates

$$d_\alpha = p_\alpha - i(P^\mu + X^{\mu'}) (\sigma_\mu\bar{\theta})_\alpha + [(\theta'\sigma^\mu\bar{\theta}) - (\theta\sigma^\mu\bar{\theta}')] (\sigma_\mu\bar{\theta})_\alpha, \quad (3.32a)$$

$$\bar{d}_{\dot{\alpha}} = \bar{p}_{\dot{\alpha}} - i(P^\mu + X^{\mu'}) (\theta\sigma_\mu)_{\dot{\alpha}} + [(\theta'\sigma^\mu\bar{\theta}) - (\theta\sigma^\mu\bar{\theta}')] (\theta\sigma_\mu)_{\dot{\alpha}}. \quad (3.32b)$$

3.2. Quantization

Since we have redefined all the fields as free fields, there is no difficulty in the quantization. We only need to replace the Dirac bracket by the quantum bracket as

$$[X^\mu(\sigma), P^\nu(\sigma')] = i\eta^{\mu\nu}\delta(\sigma - \sigma'), \quad (3.33a)$$

$$\{p_\alpha^A(\sigma), \theta^{B\beta}(\sigma')\} = -i\delta^{AB}\delta_\alpha^\beta\delta(\sigma - \sigma'), \quad (3.33b)$$

$$\{\bar{p}_{\dot{\alpha}}^A(\sigma), \bar{\theta}^{B\dot{\beta}}(\sigma')\} = -i\delta^{AB}\delta_{\dot{\alpha}}^{\dot{\beta}}\delta(\sigma - \sigma'), \quad (3.33c)$$

$$\{S(\sigma), \bar{S}(\sigma')\} = -\delta(\sigma - \sigma'). \quad (3.33d)$$

These can be translated into the OPE relations in the Euclidean formulation with radial quantization. In order to obtain the standard normalization, we set $T = 1/2\pi\alpha' = 1$ and make the replacement $P^\mu + X^{\mu'} \rightarrow 2i\partial X^\mu$, $p_\alpha \rightarrow 2ip_\alpha$, $\bar{p}_{\dot{\alpha}} \rightarrow 2i\bar{p}_{\dot{\alpha}}$, $S \rightarrow \sqrt{2}iS$ and $\bar{S} \rightarrow$

$-\sqrt{2}i\bar{S}$, where ∂ represents $\partial/\partial z$. The OPEs for the basic fields of the left (holomorphic) sector then become

$$X^\mu(z)X^\nu(w) \sim \eta^{\mu\nu} \log(z-w), \quad (3.34a)$$

$$p_\alpha(z)\theta^\beta(w) \sim \frac{\delta_\alpha^\beta}{z-w}, \quad (3.34b)$$

$$\bar{p}_{\dot{\alpha}}(z)\bar{\theta}^{\dot{\beta}}(w) \sim \frac{\delta_{\dot{\alpha}}^{\dot{\beta}}}{z-w}, \quad (3.34c)$$

$$S(z)\bar{S}(w) \sim \frac{1}{z-w}. \quad (3.34d)$$

It is also convenient to change the normalizations as

$$D_\alpha \rightarrow \frac{1}{2i}D_\alpha, \quad \bar{D}_{\dot{\alpha}} \rightarrow \frac{1}{2i}\bar{D}_{\dot{\alpha}}, \quad \Pi \rightarrow \frac{1}{2}\Pi, \quad (3.35)$$

and to rescale $\mathcal{T}$ so that $\mathcal{T} = -(1/2)\Pi^\mu\Pi_\mu/\Pi^+$.

Using these rescaled free fields, we rewrite the constraints in such a manner that separates the non-covariant fermions S and $\bar{S}$ from the remaining covariant sector. Using the super-covariant current (2.7), Π^μ can be written as

$$\Pi^+ = \pi^+, \quad (3.36a)$$

$$\Pi^- = \pi^- + \frac{1}{\pi^+} (S\partial\bar{S} - \partial S\bar{S}) - 2i\sqrt{\frac{2}{\pi^+}} (S\partial\bar{\theta}^1 + \partial\theta^1\bar{S}), \quad (3.36b)$$

$$\Pi = \pi + 2i\sqrt{\frac{2}{\pi^+}} S\partial\bar{\theta}^2, \quad (3.36c)$$

$$\bar{\Pi} = \bar{\pi} + 2i\sqrt{\frac{2}{\pi^+}} \partial\theta^2\bar{S}. \quad (3.36d)$$

The constraint generators are then rewritten in terms of the free fields as

$$D_1 = d_1 - i\sqrt{2\pi^+}\bar{S}, \quad (3.37a)$$

$$D_2 = d_2 - i\sqrt{\frac{2}{\pi^+}}\pi\bar{S} - \frac{2}{\pi^+}S\bar{S}\partial\bar{\theta}^2, \quad (3.37b)$$

$$\bar{D}_1 = \bar{d}_1 + i\sqrt{2\pi^+}S, \quad (3.37c)$$

$$\bar{D}_2 = \bar{d}_2 + i\sqrt{\frac{2}{\pi^+}}\bar{\pi}S + \frac{2}{\pi^+}S\bar{S}\partial\theta^2, \quad (3.37d)$$

$$\begin{aligned} \mathcal{T} = & -\frac{1}{2}\frac{\pi^\mu\pi_\mu}{\pi^+} - \frac{1}{2}\frac{S\partial\bar{S}}{\pi^+} + \frac{1}{2}\frac{\partial S\bar{S}}{\pi^+} + i\sqrt{\frac{2}{\pi^+}}(S\partial\bar{\theta}^1 + \partial\theta^1\bar{S}) \\ & + i\sqrt{\frac{2}{(\pi^+)^3}}\left(\bar{\pi}S\partial\bar{\theta}^2 + \pi\partial\theta^2\bar{S}\right) + 4\frac{S\bar{S}\partial\theta^2\partial\bar{\theta}^2}{(\pi^+)^2}, \end{aligned} \quad (3.37e)$$

where d_α and $\bar{d}_{\dot{\alpha}}$ are super-covariant currents (2.5).

The constraint algebras formed by the generators above are not closed in their present forms. However, these expressions are classical, although they have normal ordering ambiguities in general. What is particularly noteworthy in the $D = 10$ DS theory is that these ambiguities are used to modify the generators in such a way that their algebras become closed. A similar happens here: Including such “quantum corrections,” the constraint generators become

$$\widehat{D}_1 = D_1, \quad \widehat{\bar{D}}_{\dot{1}} = \bar{D}_{\dot{1}}, \quad (3.38a)$$

$$\widehat{D}_2 = D_2 - \frac{\partial^2 \bar{\theta}^2}{\pi^+} + \frac{1}{2} \frac{\partial \pi^+ \partial \bar{\theta}^2}{(\pi^+)^2}, \quad (3.38b)$$

$$\widehat{\bar{D}}_{\dot{2}} = \bar{D}_{\dot{2}} - \frac{\partial^2 \theta^2}{\pi^+} + \frac{1}{2} \frac{\partial \pi^+ \partial \theta^2}{(\pi^+)^2}, \quad (3.38c)$$

$$\widehat{\mathcal{T}} = \mathcal{T} + \frac{\partial \theta^2 \partial^2 \bar{\theta}^2}{(\pi^+)^2} - \frac{\partial^2 \theta^2 \partial \bar{\theta}^2}{(\pi^+)^2} - \frac{1}{8} \frac{\partial^2 \log \pi^+}{\pi^+}, \quad (3.38d)$$

and then they form a closed first-class algebra:

$$\widehat{D}_2(z) \widehat{\bar{D}}_{\dot{2}}(w) \sim \frac{4\widehat{\mathcal{T}}(w)}{z-w}, \quad (3.39)$$

with all others vanishing. Using these generators of the first-class constraints, one can straightforwardly construct the BRST charge according to the conventional procedure as

$$\tilde{Q} = \oint \frac{dz}{2\pi i} \left(\tilde{\lambda}^\alpha \widehat{D}_\alpha + \tilde{\bar{\lambda}}^{\dot{\alpha}} \widehat{\bar{D}}_{\dot{\alpha}} + c\widehat{\mathcal{T}} - 4\tilde{\lambda}^2 \tilde{\bar{\lambda}}^2 b \right), \quad (3.40)$$

where $\tilde{\lambda}^\alpha$ and $\tilde{\bar{\lambda}}^{\dot{\alpha}}$ are *unconstrained* bosonic spinor ghosts. Fermionic ghosts b and c satisfying $b(z)c(w) \sim 1/(z-w)$ are also introduced. This is exactly nilpotent quantum mechanically, due to the correction (3.38).

As in the ten-dimensional case, we can construct the composite B -ghost

$$B = b\pi^+ - \tilde{\omega}_\alpha \partial \theta^\alpha - \tilde{\bar{\omega}}_{\dot{\alpha}} \partial \bar{\theta}^{\dot{\alpha}}, \quad (3.41)$$

in terms of which the energy-momentum tensor can be expressed as

$$\begin{aligned} \{\tilde{Q}, B(z)\} &= T(z) \\ &= -\frac{1}{2} \pi^\mu \pi_\mu - \frac{1}{8} \partial^2 \log \pi^+ - \frac{1}{2} S \partial \bar{S} + \frac{1}{2} \partial S \bar{S} \\ &\quad - d_\alpha \partial \theta^\alpha - \bar{d}_{\dot{\alpha}} \partial \bar{\theta}^{\dot{\alpha}} - \tilde{\omega}_\alpha \partial \tilde{\lambda}^\alpha - \tilde{\bar{\omega}}_{\dot{\alpha}} \partial \tilde{\bar{\lambda}}^{\dot{\alpha}} - b \partial c, \end{aligned} \quad (3.42)$$

where the bosonic anti-ghosts $\tilde{\omega}_\alpha$ and $\tilde{\bar{\omega}}_{\dot{\alpha}}$ are introduced as conjugates of $\tilde{\lambda}^\alpha$ and $\tilde{\bar{\lambda}}^{\dot{\alpha}}$ satisfying

$$\tilde{\lambda}^\alpha(z)\tilde{\omega}_\beta(w) \sim \frac{\delta_\beta^\alpha}{z-w}, \quad \tilde{\bar{\lambda}}^{\dot{\alpha}}(z)\tilde{\bar{\omega}}_{\dot{\beta}}(w) \sim \frac{\delta_{\dot{\beta}}^{\dot{\alpha}}}{z-w}. \quad (3.43)$$

This energy-momentum tensor has vanishing central charge, due to the second term involving $\log \pi^+$. We can rescale the fermionic bc -ghosts to obtain the conventional reparametrization ghosts with conformal dimensions 2 and -1 as

$$\begin{aligned} e^R T e^{-R} = & -\frac{1}{2}\pi^\mu\pi_\mu + \frac{7}{8}\partial^2 \log \pi^+ - \frac{1}{2}S\partial\bar{S} + \frac{1}{2}\partial S\bar{S} \\ & - d_\alpha\partial\theta^\alpha - \bar{d}_{\dot{\alpha}}\partial\bar{\theta}^{\dot{\alpha}} - \tilde{\omega}_\alpha\partial\tilde{\lambda}^\alpha - \tilde{\bar{\omega}}_{\dot{\alpha}}\partial\tilde{\bar{\lambda}}^{\dot{\alpha}} - \partial bc - 2b\partial c \end{aligned} \quad (3.44)$$

by carrying out the similarity transformation generated by

$$R = - \oint \frac{dz}{2\pi i} bc \log \pi^+. \quad (3.45)$$

3.3. Equivalence to the PS formalism in four dimensions

Finally, we show that the cohomology of the BRST charge $\tilde{Q}$ (3.40) in the DS formalism is the same as that of the PS formalism (2.4). To begin with, we have to explicitly solve the four-dimensional pure-spinor constraint, which can also be written as

$$\lambda^\alpha \bar{\lambda}^{\dot{\alpha}} = 0. \quad (3.46)$$

This has the solutions $\lambda^\alpha = 0$ and $\bar{\lambda}^{\dot{\alpha}} = 0$.*)

As the first step to prove the equivalence, we show the decoupling of the fermionic ghost pair (b, c) and one of the two bosonic ghost pairs $(\tilde{\lambda}^2, \tilde{\omega}_2)$ and $(\tilde{\bar{\lambda}}^{\dot{2}}, \tilde{\bar{\omega}}_{\dot{2}})$. We have to consider the two cases $\tilde{\lambda}^2 \neq 0$ and $\tilde{\bar{\lambda}}^{\dot{2}} \neq 0$ separately. These correspond to the two branches of solutions of the pure-spinor constraint. The total Hilbert space is the union of these two cases.

Let us first consider the case $\tilde{\lambda}^2 \neq 0$ corresponding to the branch $\bar{\lambda}^{\dot{\alpha}} = 0$. In this case, we can consider the similarity transformation generated by

$$X = -\frac{1}{4} \oint \frac{dz}{2\pi i} \frac{c \hat{D}_{\dot{2}}}{\tilde{\lambda}^2}, \quad (3.47)$$

which transforms the BRST charge $\tilde{Q}$ into

$$e^X \tilde{Q} e^{-X} = \delta_b + Q^{(1)}, \quad (3.48)$$

*) If $\bar{\lambda}^{\dot{\alpha}}$ is the complex conjugate of λ^α , the pure-spinor constraint (3.46) has no non-trivial solution. This is a common difficulty in the PS formalism, and to avoid it we consider them to be independent fields in this section.

where

$$\delta_b = -4 \oint \frac{dz}{2\pi i} \tilde{\lambda}^2 \tilde{\lambda}^{\dot{2}} b, \quad (3.49)$$

$$Q^{(1)} = \oint \frac{dz}{2\pi i} (\tilde{\lambda}^\alpha \widehat{D}_\alpha + \tilde{\bar{\lambda}}^{\dot{1}} \widehat{D}_{\dot{1}}). \quad (3.50)$$

The general argument of homological perturbation theory⁸⁾ shows that the cohomology of $\tilde{Q}$ coincides with that of $Q^{(1)}$ in the Hilbert space without b , c , $\tilde{\lambda}^{\dot{2}}$ and $\tilde{\omega}_{\dot{2}}$ decoupled as a quartet, due to δ_b .^{*)}

We further carry out the second transformation, which consists of two successive similarity transformations. The first transformation is given by $e^Y Q^{(1)} e^{-Y}$, with

$$Y = -\frac{1}{2} \oint \frac{dz}{2\pi i} S \bar{S} \log \pi^+. \quad (3.51)$$

This yields the replacement

$$S \rightarrow \frac{S}{\sqrt{\pi^+}}, \quad \bar{S} \rightarrow \sqrt{\pi^+} \bar{S}, \quad (3.52)$$

which also implies a shift of the conformal weight of $(S, \bar{S})$ from $(1/2, 1/2)$ to $(1, 0)$. This transformation results in the decoupling of S , $\bar{S}$, $\tilde{\lambda}^{\dot{1}}$ and $\tilde{\omega}_{\dot{1}}$ as a quartet. Indeed, the BRST charge is transformed as

$$Q^{(2)} = e^Y Q^{(1)} e^{-Y} = \delta + Q + d, \quad (3.53)$$

$$\delta = \sqrt{2}i \oint \frac{dz}{2\pi i} \bar{\lambda}^{\dot{1}} S, \quad Q = \oint \frac{dz}{2\pi i} \lambda^\alpha d_\alpha, \quad (3.54)$$

$$d = \oint \frac{dz}{2\pi i} \left(-\lambda^2 \frac{\partial^2 \bar{\theta}^{\dot{2}}}{\pi^+} + \lambda^2 \frac{\partial \pi^+ \partial \bar{\theta}^{\dot{2}}}{(\pi^+)^2} + \bar{\lambda}^{\dot{1}} \bar{d}_{\dot{1}} - \sqrt{2}i \lambda^1 \pi^+ \bar{S} - \sqrt{2}i \lambda^2 \pi \bar{S} \right), \quad (3.55)$$

and the form of δ indicates such a decoupling. Subsequently, the transformation

$$e^Z Q^{(2)} e^{-Z} = \delta + Q, \quad (3.56)$$

$$\delta = \sqrt{2}i \oint \frac{dz}{2\pi i} \bar{\lambda}^{\dot{1}} S, \quad (3.57)$$

$$Q = \oint \frac{dz}{2\pi i} \lambda^\alpha d_\alpha, \quad (3.58)$$

with

$$Z = \oint \frac{dz}{2\pi i} \left(\frac{i}{\sqrt{2}} \bar{d}_{\dot{1}} \bar{S} + \frac{\partial \theta^2 \partial \bar{\theta}^{\dot{2}}}{\pi^+} \right), \quad (3.59)$$

^{*)} This is due to the fact that the cohomology of δ_b must satisfy a part of the pure-spinor constraint (3.46) $\tilde{\lambda}^2 \tilde{\lambda}^{\dot{2}} = 0$. This yields $\tilde{\lambda}^{\dot{2}} = 0$, that is, the decoupling of $\tilde{\lambda}^{\dot{2}}$, since we consider the case $\tilde{\lambda}^2 \neq 0$.

completes the second transformation. Now the final BRST charge (3.56) has the same form as the first transformation (3.48). The cohomology of $Q^{(1)}$ coincides with that of Q in the Hilbert space without S , $\bar{S}$, $\tilde{\lambda}^{\dot{1}}$ and $\tilde{\omega}_{\dot{1}}$ decoupled as a quartet, due to δ . This BRST charge Q is identical to that of the four-dimensional PS superstring in the branch $\bar{\lambda}^{\dot{\alpha}} = 0$.

We can similarly consider the other branch, $\tilde{\lambda}^{\dot{2}} \neq 0$. The first similarity transformation in this case is generated by

$$\bar{X} = -\frac{1}{4} \oint \frac{dz}{2\pi i} \frac{c\hat{D}_2}{\bar{\lambda}^{\dot{2}}}, \quad (3.60)$$

which transforms the BRST charge as

$$e^{\bar{X}} \tilde{Q} e^{-\bar{X}} = \delta_b + Q^{(1)}, \quad (3.61)$$

$$\delta_b = -4 \oint \frac{dz}{2\pi i} \lambda^2 \bar{\lambda}^{\dot{2}} b, \quad (3.62)$$

$$Q^{(1)} = \oint \frac{dz}{2\pi i} (\lambda^1 \hat{D}_1 + \bar{\lambda}^{\dot{\alpha}} \hat{D}_{\dot{\alpha}}). \quad (3.63)$$

Then λ^2 , ω_2 , c and b are decoupled as a quartet.

The second similarity transformation, given by

$$\bar{Y} = \frac{1}{2} \oint \frac{dz}{2\pi i} S \bar{S} \log \pi^+, \quad (3.64a)$$

$$\bar{Z} = - \oint \frac{dz}{2\pi i} \left(\frac{i}{\sqrt{2}} d_1 S + \frac{\partial \theta^2 \partial \bar{\theta}^{\dot{2}}}{\pi^+} \right), \quad (3.64b)$$

leads to

$$e^{\bar{Z}} e^{\bar{Y}} Q^{(1)} e^{-\bar{Y}} e^{-\bar{Z}} = \delta + Q, \quad (3.65)$$

$$\delta = -\sqrt{2}i \oint \frac{dz}{2\pi i} \lambda^1 \bar{S}, \quad (3.66)$$

$$Q = \oint \frac{dz}{2\pi i} \bar{\lambda}^{\dot{\alpha}} \bar{d}_{\dot{\alpha}}, \quad (3.67)$$

and λ^1 , ω_1 , S and $\bar{S}$ are decoupled as a quartet. The BRST charge Q is that of the four-dimensional PS superstring in the branch $\lambda^{\alpha} = 0$.

Thus the physical states are finally obtained as the union of the cohomologies of the BRST charges (3.58) and (3.67). They coincide with those of the four-dimensional PS superstring defined by the cohomology of the BRST charge with the constrained spinor (2.4).

§4. DS formalism in six dimensions

Similarly to the previous section, in this section we consider the six-dimensional superstring in the DS formalism.

4.1. Lagrangian, symmetries and constraints

The six-dimensional superstring in the DS formalism is defined using superspace coordinates $(x^\mu, \theta_I^{A\alpha})$ and the additional fermionic field $\tilde{\theta}_I^{A\alpha}$ ($A = 1, 2$). Using these fundamental fields, the Lagrangian is given by $\mathcal{L} = \mathcal{L}_K + \mathcal{L}_{WZ}$, with

$$\mathcal{L}_K = -\frac{1}{2}\sqrt{-g}g^{ab}\Pi_a^\mu\Pi_{\mu b}, \quad (4.1a)$$

$$\mathcal{L}_{WZ} = \epsilon^{ab}\Pi_a^\mu(W_{\mu b} - \hat{W}_{\mu b}) - \epsilon^{ab}W_a^\mu\hat{W}_{\mu b}, \quad (4.1b)$$

where

$$\Pi_a^\mu = \partial_a X^\mu - \sum_{A=1}^2 i\partial_a(\theta^{IA}C\gamma^\mu\tilde{\theta}_I^A) - \sum_{A=1}^2 W_a^{A\mu}, \quad (4.2)$$

$$W_a^{A\mu} = i(\Theta^{IA}C\gamma^\mu\partial_a\Theta_I^A), \quad (4.3)$$

$$\Theta_I^A = \tilde{\theta}_I^A - \theta_I^A. \quad (4.4)$$

This Lagrangian is invariant under the world-sheet reparametrization and the space-time Poincaré transformation, where the space-time supersymmetry is defined by

$$\delta\theta_I^A = \epsilon_I^A, \quad \delta\tilde{\theta}_I^A = 0, \quad (4.5a)$$

$$\delta X^\mu = \sum_{A=1}^2 i(\epsilon^{IA}C\gamma^\mu\theta_I^A). \quad (4.5b)$$

In six dimensions, the local supersymmetry is given by

$$\delta\theta_I^A = \chi_I^A, \quad \delta\tilde{\theta}_I^A = \chi_I^A, \quad (4.6a)$$

$$\delta X^\mu = \sum_{A=1}^2 i(\chi^{IA}C\gamma^\mu\Theta_I^A), \quad (4.6b)$$

which guarantees the equivalence to the conventional Green-Schwarz formalism.

The canonical conjugate of X^μ is obtained as

$$k_\mu = -\sqrt{-g}g^{0b}\Pi_{\mu b} + W_{\mu 1} - \hat{W}_{\mu 1}. \quad (4.7)$$

By computing the canonical conjugates k_α^{IA} and $\tilde{k}_\alpha^{IA}$ of the fermionic fields $\theta_I^{\alpha A}$ and $\tilde{\theta}_I^{\alpha A}$, we obtain the primary constraints

$$D_\alpha^{IA} = p_\alpha^{IA} - iP^\mu(C\gamma_\mu\tilde{\theta}^{IA})_\alpha - i(P^\mu + \eta^A(\Pi_1^\mu + W_1^{\mu A'}))(C\gamma_\mu\Theta^{IA})_\alpha \approx 0, \quad (4.8a)$$

$$\tilde{D}_\alpha^{IA} = \tilde{p}_\alpha^{IA} + iP^\mu(C\gamma_\mu\theta^{IA})_\alpha + i(P^\mu + \eta^A(\Pi_1^\mu + W_1^{\mu A'}))(C\gamma_\mu\Theta^{IA})_\alpha \approx 0, \quad (4.8b)$$

where $\eta_1 = -\eta_2 = 1$ and $A' = 2(1)$ for $A = 1(2)$. We also define the generators $\Delta_\alpha^{IA} = D_\alpha^{IA} + \tilde{D}_\alpha^{IA}$.

Using the same ADM decomposition of the world-sheet metric as in the four-dimensional case (3.9), we can obtain the Hamiltonian as

$$\begin{aligned}\mathcal{H} &= \frac{N}{\sqrt{\gamma}} \frac{1}{2} \left((P_\mu - W_{\mu 1} + \hat{W}_{\mu 1})(P^\mu - W_1^\mu + \hat{W}_1^\mu) + \Pi_1^\mu \Pi_{\mu 1} \right) \\ &\quad + N^1 (P_\mu - W_{\mu 1} + \hat{W}_{\mu 1}) \Pi_1^\mu + \sum_A \dot{\theta}_I^A D^{IA} + \sum_A \dot{\tilde{\theta}}_I^A \tilde{D}^{IA}, \\ &= \frac{N}{\sqrt{\gamma}} T_0 + N^1 T_1 + \sum_A \dot{\theta}_I^A D^{IA} + \sum_A \dot{\tilde{\theta}}_I^A \tilde{D}^{IA}.\end{aligned}\tag{4.9}$$

Here, the energy-momentum tensors are given by

$$T_+ = \frac{1}{2}(T_0 + T_1) = \frac{1}{4} \Pi^\mu \Pi_\mu, \tag{4.10a}$$

$$T_- = \frac{1}{2}(T_0 - T_1) = \frac{1}{4} \hat{\Pi}^\mu \hat{\Pi}_\mu, \tag{4.10b}$$

with

$$\begin{aligned}\Pi^\mu &= k^\mu - W_1^\mu + \hat{W}_1^\mu + \Pi_1^\mu, \\ &= k^\mu + X'^\mu - \sum_A i(\theta^{IA} C \gamma^\mu \tilde{\theta}_I^A)' - 2W_1^\mu,\end{aligned}\tag{4.11a}$$

$$\begin{aligned}\hat{\Pi}^\mu &= k^\mu - W_1^\mu + \hat{W}_1^\mu - \Pi_1^\mu, \\ &= k^\mu - X'^\mu + \sum_A i(\theta^{IA} C \gamma^\mu \tilde{\theta}_I^A)' + 2\hat{W}_1^\mu.\end{aligned}\tag{4.11b}$$

Then we can set the canonical Poisson brackets as

$$\{X^\mu(\sigma), P^\nu(\sigma')\}_P = \eta^{\mu\nu} \delta(\sigma - \sigma'), \tag{4.12a}$$

$$\{\theta_I^{A\alpha}(\sigma), p_\beta^{JB}(\sigma')\}_P = -\delta^{AB} \delta_I^J \delta_\beta^\alpha \delta(\sigma - \sigma'), \tag{4.12b}$$

$$\{\tilde{\theta}_I^{A\alpha}(\sigma), \tilde{p}_\beta^{JB}(\sigma')\}_P = -\delta^{AB} \delta_I^J \delta_\beta^\alpha \delta(\sigma - \sigma'). \tag{4.12c}$$

Using these canonical Poisson brackets, we can calculate the constraint algebra of the fermionic generators $(\Delta_\alpha^I, \tilde{D}_\alpha^I)$ as

$$\{\tilde{D}_\alpha^I(\sigma), \tilde{D}_\beta^J(\sigma')\}_P = -2i\epsilon^{IJ}(C\gamma_\mu)_{\alpha\beta} \Pi^\mu(\sigma) \delta(\sigma - \sigma'), \tag{4.13}$$

with all others vanishing.

The total energy-momentum tensor is now given by

$$T = T_+ + t_+ = \frac{1}{4} \Pi^\mu \Pi_\mu + \Theta_I'^\alpha \tilde{D}_\alpha^I, \tag{4.14}$$

which forms the Virasoro algebra:

$$\{T_+(\sigma), T_+(\sigma')\}_P = 2T_+(\sigma)\delta'(\sigma - \sigma') + T'(\sigma)\delta(\sigma - \sigma'). \quad (4.15)$$

We can decompose the constraint $\tilde{D}_\alpha^I \approx 0$ into the second-class constraint $\tilde{D}_a^I \approx 0$ and the first-class constraint $K_a^I \approx 0$ as

$$K_a^I = \tilde{D}_a^I + \frac{(\epsilon \mathbb{I})_{\dot{a}b}}{\Pi^+} \tilde{D}^{Ib}. \quad (4.16)$$

These $\tilde{D}_a^I$ satisfy

$$\{\tilde{D}_a^I(\sigma), \tilde{D}_b^J(\sigma')\}_P = -2i\epsilon^{IJ}\epsilon_{ab}\Pi^+(\sigma)\delta(\sigma - \sigma'), \quad (4.17a)$$

$$\{K_a^I(\sigma), K_b^J(\sigma')\}_P = -8i\epsilon^{IJ}\epsilon_{\dot{a}b}(\mathcal{T} + \mathcal{K})(\sigma)\delta'(\sigma - \sigma') + \tilde{D}\text{-terms}, \quad (4.17b)$$

where $(\epsilon \mathbb{I})_{\dot{a}b} = (\epsilon \gamma_i)_{\dot{a}b}\Pi^i$, $\mathcal{T} = \frac{T}{\Pi^+}$ and $\mathcal{K} = \frac{K_\epsilon^K \Theta_K^\epsilon}{\Pi^+}$. The $\tilde{D}$ -terms in (4.17b) can be set equal to zero after taking the Dirac bracket.

By choosing the semi-light-cone gauge $\tilde{\theta}_I^{\dot{a}} \approx 0$ for the first-class constraint, we finally have three second-class constraints:

$$\tilde{D}_a^I \approx 0, \quad K_a^I \approx 0, \quad \tilde{\theta}_i^{\dot{a}} \approx 0. \quad (4.18)$$

The Dirac bracket for these second-class constraints is defined by

$$\begin{aligned} \{A(\sigma), B(\sigma')\}_D = & \{A(\sigma), B(\sigma')\}_P \\ & + \int d\sigma_1 \{A(\sigma), \tilde{D}_a^I(\sigma_1)\}_P \frac{\epsilon_{IJ}\epsilon^{ab}}{2i\Pi^+}(\sigma_1) \{\tilde{D}_b^J(\sigma_1), B(\sigma')\}_P \\ & + \int d\sigma_1 \{A(\sigma), K_a^I(\sigma_1)\}_P \{\tilde{\theta}_I^{\dot{a}}(\sigma_1), B(\sigma')\}_P \\ & + \int d\sigma_1 \{A(\sigma), \tilde{\theta}_I^{\dot{a}}(\sigma_1)\}_P \{K_a^I(\sigma_1), B(\sigma')\}_P \\ & - 8i \int d\sigma_1 \{A(\sigma), \tilde{\theta}_I^{\dot{a}}(\sigma_1)\}_P \epsilon_{\dot{a}b}^{IJ} \mathcal{T}(\sigma_1) \{\tilde{\theta}_J^{\dot{b}}(\sigma_1), B(\sigma')\}_P. \end{aligned} \quad (4.19)$$

In this semi-light cone gauge, the independent fields are $(X^\mu, P_\mu, \theta_I^\alpha, p_\alpha^I, \tilde{\theta}_I^a)$, which on the constraint plane $\tilde{\theta}_I^{\dot{a}} \approx 0$ satisfy, for example,

$$\{\tilde{\theta}_I^a(\sigma), \tilde{\theta}_J^b(\sigma')\}_D = -\frac{i\epsilon_{IJ}\epsilon^{ab}}{2\Pi^+}(\sigma)\delta(\sigma - \sigma'), \quad (4.20a)$$

$$\{X^-(\sigma), \tilde{\theta}_I^a(\sigma')\}_D = -\frac{1}{\Pi^+}\tilde{\theta}_I^a(\sigma)\delta(\sigma - \sigma'), \quad (4.20b)$$

$$\{X^-(\sigma), P^-(\sigma')\}_D = \frac{2i}{\Pi^+}\tilde{\theta}_I^a\theta_a^I(\sigma)\delta'(\sigma - \sigma'), \quad (4.20c)$$

$$\{X^-(\sigma), P^i(\sigma')\}_D = -\frac{i}{\Pi^+} \tilde{\theta}_I^a(\epsilon \bar{\gamma}^i)_{ab} \theta^{Ib}(\sigma) \delta'(\sigma - \sigma'), \quad (4.20d)$$

$$\{\tilde{\theta}_I^a(\sigma), P^-(\sigma')\}_D = -\frac{1}{\Pi^+} \Theta_I^a(\sigma) \delta'(\sigma - \sigma'), \quad (4.20e)$$

$$\{\tilde{\theta}_I^a(\sigma), P^i(\sigma')\}_D = \frac{1}{2\Pi^+} (\bar{\gamma}^i)^a_{\dot{c}} \theta_I^{\dot{c}}(\sigma) \delta(\sigma - \sigma'). \quad (4.20f)$$

We can again redefine the fields so that the new ones become free. We first redefine the non-covariant fermionic field as

$$S_I^a = \sqrt{2\Pi^+} \tilde{\theta}_I^a. \quad (4.21)$$

Then S_I^a satisfies the free relations

$$\{S_I^a(\sigma), S_J^b(\sigma')\}_D = -i\epsilon_{IJ} \epsilon^{ab} \delta(\sigma - \sigma'), \quad (4.22a)$$

$$\{X^-(\sigma), S_I^a(\sigma')\}_D = 0. \quad (4.22b)$$

Similarly, we can obtain the free fields through the redefinitions

$$P^\mu = k^\mu - i(\tilde{\theta}^K C \gamma^\mu \theta_K)' + i(\hat{\tilde{\theta}}^K C \gamma^\mu \hat{\theta}_K)', \quad (4.23a)$$

$$p_a^{IA} = k_a^{IA} + \eta^A \left(-iX'^+ \tilde{\theta}_a^{IA} - 4\theta_c^{KA} \theta'^{\dot{c}A} \tilde{\theta}_a^{IA} + 2\theta_c^{IA} (\theta_K^{\dot{c}A} \tilde{\theta}_a^{KA})' + 2(\theta_c^{IA} \theta_K^{\dot{c}A})' \tilde{\theta}_a^{KA} \right), \quad (4.23b)$$

$$p_{\dot{a}}^{IA} = k_{\dot{a}}^{IA} + \eta^A \left(i(\epsilon X')_{\dot{a}c} \tilde{\theta}^{IcA} - 2\theta_{\dot{a}}^{IA} \theta_c^{KA} \tilde{\theta}_K^{cA} - 2\theta_{\dot{a}}^{KA} \theta_c^{IA} \tilde{\theta}_K^{cA} - 6\theta_{\dot{a}}^{IA} \theta_c^{KA} \tilde{\theta}_K^{cA} \right. \\ \left. - 6\theta_{\dot{a}}^{KA} \theta_c^{IA} \tilde{\theta}_K^{cA} - 2\theta_{\dot{a}}^{IA} \tilde{\theta}_c^{KA} \tilde{\theta}_K^{cA} + 4\theta_{\dot{a}}^{KA} \tilde{\theta}_c^{IA} \tilde{\theta}_K^{cA} - 2(\theta_{\dot{a}}^{KA} \tilde{\theta}_{Kc}^{A'})' \tilde{\theta}^{IcA} \right). \quad (4.23c)$$

These redefined fields satisfy

$$\{X^\mu(\sigma), P^\nu(\sigma')\}_D = \eta^{\mu\nu} \delta(\sigma - \sigma'), \quad (4.24a)$$

$$\{\theta_I^{A\alpha}(\sigma), p_\beta^{BJ}(\sigma')\}_D = -\delta^{AB} \delta_I^J \delta_\beta^\alpha \delta(\sigma - \sigma'), \quad (4.24b)$$

with the remaining brackets vanishing.

4.2. Quantization

The quantization is obtained by replacing the Dirac bracket with the (anti-)commutation relations, which can be rewritten in the form of the operator product expansion in the conventional radial quantization, after the appropriate field rescalings:

$$X^\mu(z) X^\nu(w) \sim \eta^{\mu\nu} \log(z - w), \quad (4.25a)$$

$$p_\alpha^I(z) \theta_J^\beta(w) \sim \frac{\delta_J^I \delta_\alpha^\beta}{z - w}, \quad (4.25b)$$

$$S_I^a(z) S_J^b(w) \sim -\frac{\epsilon_{IJ} \epsilon^{ab}}{z - w}. \quad (4.25c)$$

The constraint generators are classically given by

$$D_a^I = d_a^I + \sqrt{2\pi^+} S_a^I, \quad (4.26a)$$

$$\begin{aligned} D_{\dot{a}}^I &= d_{\dot{a}}^I - \sqrt{\frac{2}{\pi^+}} (\epsilon \not{\pi})_{\dot{a}b} S^{Ib} + \frac{2}{\pi^+} S_c^I S_K^c \partial \theta_{\dot{a}}^K, \\ &= d_{\dot{a}}^I - \sqrt{\frac{2}{\pi^+}} (\epsilon \not{\pi})_{\dot{a}b} S^{Ib} - \frac{1}{\pi^+} (\epsilon \gamma_i)_{\dot{a}b} S^{Ib} S^{Kc} (\epsilon \bar{\gamma}_i)_{cd} \partial \theta_K^d, \end{aligned} \quad (4.26b)$$

$$\begin{aligned} \mathcal{T} &= -\frac{1}{2} \frac{\pi^\mu \pi_\mu}{\pi^+} - \frac{1}{2} \frac{S_a^K \partial S_K^a}{\pi^+} - \sqrt{\frac{2}{\pi^+}} \partial \theta_a^K S_K^a \\ &\quad - \sqrt{\frac{2}{\pi^+}} \frac{\partial \theta^{K\dot{a}} (\epsilon \not{\pi})_{\dot{a}b} S_K^b}{\pi^+} - 2 \frac{\partial \theta^{K\dot{a}} \partial \theta_{\dot{a}}^L S_K^b S_{Lb}}{(\pi^+)^2}, \end{aligned} \quad (4.26c)$$

where the super-covariant currents d_a^I and π^μ are defined by (2.15). Including “quantum” corrections, as we have done in $D = 4$, they become^{*)}

$$\widehat{D}_a^I = D_a^I, \quad (4.27a)$$

$$\widehat{D}_{\dot{a}}^I = D_{\dot{a}}^I - 2 \frac{\partial^2 \theta_{\dot{a}}^I}{\pi^+} + \frac{\partial \pi^+ \partial \theta_{\dot{a}}^I}{(\pi^+)^2} + \frac{8}{3} \frac{\partial \theta_{\dot{c}}^I \partial \theta_K^{\dot{c}} \partial \theta_{\dot{a}}^K}{(\pi^+)^2}, \quad (4.27b)$$

$$\widehat{\mathcal{T}} = \mathcal{T} - \frac{1}{4} \frac{\partial^2 \log \pi^+}{\pi^+} - 2 \frac{\partial^2 \theta_{\dot{c}}^K \partial \theta_K^{\dot{c}}}{(\pi^+)^2} - \frac{8}{3} \frac{\partial \theta_{\dot{c}}^K \partial \theta_K^{\dot{c}} \partial \theta_{\dot{a}}^L \partial \theta_L^{\dot{a}}}{(\pi^+)^3}, \quad (4.27c)$$

and satisfy

$$\widehat{D}_{\dot{a}}^I(z) \widehat{D}_{\dot{b}}^J(w) \sim - \frac{4 \epsilon^{IJ} \epsilon_{\dot{a}\dot{b}} \widehat{\mathcal{T}}(w)}{z - w}, \quad (4.28a)$$

$$\text{rest} \sim 0. \quad (4.28b)$$

The BRST charge is straightforwardly constructed from this constraint algebra as

$$\tilde{Q} = \oint \frac{dz}{2\pi i} \left(\tilde{\lambda}_I^\alpha \widehat{D}_\alpha^I + c \widehat{\mathcal{T}} - 2 \tilde{\lambda}_a^I \tilde{\lambda}_{\dot{a}}^{\dot{a}} b \right), \quad (4.29)$$

with the unconstrained bosonic ghost $\tilde{\lambda}_I^\alpha$ and the fermionic ghost pair b and c with $b(z)c(w) \sim 1/(z - w)$. This BRST charge (4.29) is exactly nilpotent.

We can define the composite B -field as

$$B = b\pi^+ - \tilde{\omega}_\alpha^I \partial \theta_I^\alpha, \quad (4.30)$$

^{*)} We should note that, in contrast to the $D = 4$ case, the extra terms in Eqs. (4.27b) and (4.27c) *cannot* be regarded as coming solely from the normal ordering ambiguities; the origin of these terms is unclear. In any case, however, the addition of these terms closes the algebra and leads to a consistent theory, as we show below.

where $\tilde{\omega}_\alpha^I$ is the conjugate bosonic anti-ghost satisfying

$$\tilde{\lambda}_I^\alpha(z)\tilde{\omega}_\beta^J(w) \sim \frac{\delta_I^J \delta_\beta^\alpha}{z-w}. \quad (4.31)$$

Then, the energy-momentum tensor can be obtained as

$$\begin{aligned} \{\tilde{Q}, B(z)\} &= T(z), \\ T &= -\frac{1}{2}\pi^\mu\pi_\mu - \frac{1}{4}\partial^2 \log \pi^+ - \frac{1}{2}S_a^I \partial S_I^a - d_\alpha^I \partial \theta_I^\alpha - \omega_\alpha^I \partial \lambda_I^\alpha - b\partial c. \end{aligned} \quad (4.32)$$

This energy-momentum tensor has vanishing central charge, and the similarity transformation defined by

$$R = - \oint \frac{dz}{2\pi i} bc \log \pi^+ \quad (4.33)$$

yields the conventional one with the reparametrization ghosts

$$e^R T e^{-R} = -\frac{1}{2}\pi^\mu\pi_\mu + \frac{3}{4}\partial^2 \log \pi^+ - \frac{1}{2}S_a^I \partial S_I^a - d_\alpha^I \partial \theta_I^\alpha - \omega_\alpha^I \partial \lambda_I^\alpha - \partial bc - 2b\partial c. \quad (4.34)$$

4.3. Equivalence to the PS formalism in six dimensions

Before showing coincidence of the physical spectra of the DS and PS formalisms, we have to explicitly solve the pure-spinor constraint in six dimensions. The pure-spinor constraint in six dimensions can be written

$$\lambda_1^\alpha (C\gamma^\mu)_{\alpha\beta} \lambda_2^\beta = 0, \quad (4.35)$$

where $\lambda_{1,2}^\alpha = \lambda_{I=1,2}^\alpha$. This equation can be satisfied if λ_2^α is proportional to λ_1^α .^{*)} Because the proportionality constant is arbitrary, the number of independent degrees of freedom of a pure spinor λ_I^α becomes $8 - 3 = 5$. Equation (4.35) can be explicitly solved by using the light-cone decomposition as follows. Let us first consider a component of (4.35),

$$\lambda_1 C\gamma^+ \lambda_2 = \lambda_{1\dot{a}} \lambda_2^{\dot{a}} = 0. \quad (4.36)$$

If we introduce the spinor dual to $\lambda_1^{\dot{a}}$ as $\lambda_{1\dot{a}} l^{\dot{a}} = 1$, the arbitrary spinor can be expanded in $\lambda_1^{\dot{a}}$ and $l^{\dot{a}}$ as

$$\lambda_2^{\dot{a}} = \alpha \lambda_1^{\dot{a}} + \tilde{\alpha} l^{\dot{a}}, \quad (4.37)$$

where the expansion coefficients are

$$\alpha = \lambda_{2\dot{a}} l^{\dot{a}}, \quad \tilde{\alpha} = -\lambda_{2\dot{a}} \lambda_1^{\dot{a}}. \quad (4.38)$$

^{*)} As in the four-dimensional case, we consider λ_1^α and λ_2^α to be two independent Weyl spinors, which are actually complex conjugates, due to the $SU(2)$ -Majorana constraint (A.21b).

Because the constraint (4.36) leads to $\tilde{\alpha} = 0$, we have

$$\lambda_2^{\dot{a}} = \alpha \lambda_1^{\dot{a}} \quad (4.39)$$

for the pure spinor. The next constraint,

$$\begin{aligned} \lambda_1 C \gamma^i \lambda_2 &= \lambda_1^{\dot{a}} (\epsilon \tilde{\gamma}^i)_{\dot{a}b} \lambda_2^b - \lambda_2^{\dot{b}} (\epsilon \tilde{\gamma}^i)_{\dot{b}a} \lambda_1^a \\ &= \lambda_1^{\dot{a}} (\epsilon \tilde{\gamma}^i)_{\dot{a}b} (\lambda_2^b - \alpha \lambda_1^b) = 0, \end{aligned} \quad (4.40)$$

is satisfied by

$$\lambda_2^a = \alpha \lambda_1^a, \quad (4.41)$$

with which the final constraint, $\lambda_1 C \gamma^- \lambda_2 = 0$, is automatically satisfied. In summary, we can satisfy the pure-spinor constraint with $\lambda_2^a = \alpha \lambda_1^a$, where $\alpha = \lambda_{2\dot{a}} l^{\dot{a}}$ with $\lambda_{1\dot{a}} l^{\dot{a}} = 1$.

Now we can show that the cohomology of the BRST charge $\tilde{Q}$ (4.29) of the DS formalism coincides with that of the PS formalism (2.13). Using the first similarity transformation generated by

$$X = \frac{1}{4} \oint \frac{dz}{2\pi i} c (l^{\dot{a}} \hat{D}_{\dot{a}}^2), \quad (4.42)$$

the BRST charge becomes

$$e^X \tilde{Q} e^{-X} = \delta_b + Q^{(1)}, \quad (4.43)$$

$$\delta_b = - \oint \frac{dz}{2\pi i} 4\tilde{\alpha} b, \quad (4.44)$$

$$Q^{(1)} = \oint \frac{dz}{2\pi i} (\tilde{\lambda}_I^a \hat{D}_I^a + \tilde{\lambda}_1^{\dot{a}} (\hat{D}_{\dot{a}}^1 + \alpha \hat{D}_{\dot{a}}^2)), \quad (4.45)$$

where $\alpha = \tilde{\lambda}_{2\dot{a}} l^{\dot{a}}$ and $\tilde{\alpha} = -\tilde{\lambda}_{2\dot{a}} \tilde{\lambda}_1^{\dot{a}}$. Then $\tilde{\alpha}, \tilde{\beta}, c$ and b are decoupled as a quartet, where $\tilde{\beta}$ is the conjugate of $\tilde{\alpha}$ which can, in principle, be constructed from $\tilde{\omega}$.

The second similarity transformations, generated by

$$Y = - \frac{1}{2} \oint \frac{dz}{2\pi i} S_a^1 S^{2a} \log \pi^+, \quad (4.46a)$$

$$Z^1 = \oint \frac{dz}{2\pi i} \left(\frac{1}{\sqrt{2}} \left(S^{1a} + \frac{\alpha}{\pi^+} S^{2a} \right) d_a^2 + \frac{2}{\alpha} \frac{\partial \theta^{1\dot{a}}}{\pi^+} (\partial \theta_{\dot{a}}^1 + \alpha \partial \theta_{\dot{a}}^2) \right), \quad (4.46b)$$

$$Z^2 = \frac{\alpha}{2} \oint \frac{dz}{2\pi i} \frac{S_a^2 S^{2a}}{\pi^+}, \quad (4.46c)$$

give

$$e^{Z^2} e^{Z^1} e^Y Q^{(1)} e^{-Y} e^{-Z^1} e^{-Z^2} = \delta + Q, \quad (4.47)$$

$$\delta = \oint \frac{dz}{2\pi i} \sqrt{2} (\tilde{\lambda}_2^a - \alpha \tilde{\lambda}_1^a) S_a^2, \quad (4.48)$$

$$Q = \oint \frac{dz}{2\pi i} \left(\tilde{\lambda}_1^a (d_a^1 + \alpha d_a^2) + \tilde{\lambda}_1^{\dot{a}} (d_{\dot{a}}^1 + \alpha d_{\dot{a}}^2) \right). \quad (4.49)$$

Then $(\tilde{\lambda}_2^a - \alpha \tilde{\lambda}_1^a)$, $(\tilde{\omega}_a^2 - \alpha \tilde{\omega}_a^1)$, S_a^1 and S_a^2 are decoupled as a quartet. The final form of Q (4.49) can be written as

$$Q = \oint \frac{dz}{2\pi i} \tilde{\lambda}_I^\alpha d_\alpha^I, \quad (4.50)$$

with the constraint $\tilde{\lambda}_{2\alpha} = \alpha \tilde{\lambda}_{1\alpha}$, which is the BRST charge of the PS formalism (2.13).

§5. Coupling to the Calabi-Yau sector

In the conventional formulations, a lower-dimensional superstring is not consistent by itself but must be combined with additional degrees of freedom which come from the compactified space. Here, since we assume the lower-dimensional supersymmetry, it is known that such degrees of freedom are represented by some unitary representations of $N = 2$ superconformal field theory, which we describe by using the generators $(T_C, G_C^\pm, J_C)$ satisfying

$$T_C(z)T_C(w) \sim \frac{c/2}{(z-w)^4} + \frac{2T_C(w)}{(z-w)^2} + \frac{\partial T_C(w)}{z-w}, \quad (5.1a)$$

$$T_C(z)G_C^\pm(w) \sim \frac{\frac{3}{2}G_C^\pm(w)}{(z-w)^2} + \frac{\partial G_C^\pm(w)}{z-w}, \quad T_C(z)J_C(w) \sim \frac{J_C(w)}{(z-w)^2} + \frac{\partial J_C(w)}{z-w}, \quad (5.1b)$$

$$G_C^+(z)G_C^-(w) \sim \frac{c/3}{(z-w)^3} + \frac{J_C(w)}{(z-w)^2} + \frac{T_C(w) + \frac{1}{2}\partial J_C(w)}{z-w}, \quad (5.1c)$$

$$J_C(z)G_C^\pm(w) \sim \pm \frac{G_C^\pm(w)}{z-w}, \quad J_C(z)J_C(w) \sim \frac{c/3}{(z-w)^2}, \quad (5.1d)$$

where $c = 9$ for $d = 4$ and $c = 6$ for $d = 6$. We call this the Calabi-Yau (CY) sector.

In order to combine the lower-dimensional superstring with this CY sector, the hidden topological $N = 2$ superconformal symmetry⁹⁾⁻¹¹⁾ generated by

$$G^+ = J_{\text{BRST}} = \tilde{\lambda}^\alpha \widehat{D}_\alpha + \tilde{\tilde{\lambda}}^{\dot{\alpha}} \widehat{\bar{D}}_{\dot{\alpha}} + c\widehat{\mathcal{T}} - 4\tilde{\lambda}^2 \tilde{\tilde{\lambda}}^{\dot{2}} b, \quad (5.2a)$$

$$G^- = B = b\pi^+ - \tilde{\omega}_\alpha \partial \theta^\alpha - \tilde{\tilde{\omega}}_{\dot{\alpha}} \partial \bar{\theta}^{\dot{\alpha}}, \quad (5.2b)$$

$$\begin{aligned} T = & -\frac{1}{2}\pi^\mu \pi_\mu - \frac{1}{8}\partial^2 \log \pi^+ - \frac{1}{2}S\partial \bar{S} + \frac{1}{2}\partial S \bar{S} \\ & - d_\alpha \partial \theta^\alpha - \bar{d}_{\dot{\alpha}} \partial \bar{\theta}^{\dot{\alpha}} - \tilde{\omega}_\alpha \partial \tilde{\lambda}^\alpha - \tilde{\tilde{\omega}}_{\dot{\alpha}} \partial \tilde{\tilde{\lambda}}^{\dot{\alpha}} - b\partial c, \end{aligned} \quad (5.2c)$$

$$J = bc - \tilde{\lambda}^\alpha \tilde{\omega}_\alpha - \tilde{\tilde{\lambda}}^{\dot{\alpha}} \tilde{\tilde{\omega}}_{\dot{\alpha}}, \quad (5.2d)$$

for $d = 4$, and

$$G^+ = J_{\text{BRST}} = \tilde{\lambda}_I^\alpha \widehat{D}_\alpha^I + c\widehat{\mathcal{T}} - 2\tilde{\lambda}_a^I \tilde{\tilde{\lambda}}_I^{\dot{a}} b, \quad (5.3a)$$

$$G^- = B = b\pi^+ - \tilde{\omega}_\alpha^I \partial \theta_I^\alpha, \quad (5.3b)$$

$$T = -\frac{1}{2}\pi^\mu\pi_\mu - \frac{1}{4}\partial^2\log\pi^+ - \frac{1}{2}S_a^I\partial S_I^a - d_\alpha^I\partial\theta_I^\alpha - \tilde{\omega}_\alpha^I\partial\tilde{\lambda}_I^\alpha - b\partial c, \quad (5.3c)$$

$$J = bc - \tilde{\lambda}_I^\alpha\tilde{\omega}_\alpha^I, \quad (5.3d)$$

for $d = 6$, play an important role. We can couple this topological superconformal algebra (SCA) with the $N = 2$ SCA in the CY sector by twisting it so as to obtain the topological SCA. This is consistent with the fact that the central charge of the lower-dimensional superstring is already zero, as explained above. The energy-momentum tensor in the CY sector must also have vanishing central charge. The BRST charge of the coupled system is then given by

$$Q_B = \oint \frac{dz}{2\pi i} (G^+ + G_C^+). \quad (5.4)$$

After applying the same similarity transformations as in the case without the CY sector, the cohomology is equivalent to that of the PS formalism coupled to the topological string in the Calabi-Yau space with the BRST charge

$$Q = Q_{PS} + \oint \frac{dz}{2\pi i} G_C^+, \quad (5.5)$$

where Q_{PS} is given by (2.4) or (2.13). The four-dimensional case is treated in 12).

Another way to couple the lower-dimensional superstring to the CY sector is by extending our constraint generators to those forming the same algebra but including the generators of $N = 2$ SCA in the CY sector. This is possible in the $d = 4$ case if we choose

$$\hat{D}_1 = D_1, \quad \hat{\bar{D}}_1 = \bar{D}_1, \quad (5.6a)$$

$$\hat{D}_2 = D_2 + \frac{2}{\sqrt{\pi^+}}G_C^+ + 2\frac{\partial\bar{\theta}^2 J_C}{\pi^+} - 4\frac{\partial^2\bar{\theta}^2}{\pi^+} + 2\frac{\partial\pi^+\partial\bar{\theta}^2}{(\pi^+)^2}, \quad (5.6b)$$

$$\hat{\bar{D}}_2 = \bar{D}_2 + \frac{2}{\sqrt{\pi^+}}G_C^- - 2\frac{\partial\theta^2 J_C}{\pi^+} - 4\frac{\partial^2\theta^2}{\pi^+} + 2\frac{\partial\pi^+\partial\theta^2}{(\pi^+)^2}, \quad (5.6c)$$

$$\hat{T} = T + \frac{T_C}{\pi^+} + 4\frac{\partial\theta^2\partial^2\bar{\theta}^2}{(\pi^+)^2} - 4\frac{\partial^2\theta^2\partial\bar{\theta}^2}{(\pi^+)^2} - \frac{1}{2}\frac{\partial^2\log\pi^+}{\pi^+}. \quad (5.6d)$$

The energy-momentum tensor then becomes

$$\begin{aligned} \{\tilde{Q}, B(z)\} &= T(z), \\ T &= -\frac{1}{2}\pi^\mu\pi_\mu - \frac{1}{2}\partial^2\log\pi^+ - \frac{1}{2}S\partial\bar{S} + \frac{1}{2}\partial S\bar{S} \\ &\quad - d_\alpha\partial\theta^\alpha - \bar{d}_{\dot{\alpha}}\partial\bar{\theta}^{\dot{\alpha}} - \omega_\alpha\partial\lambda^\alpha - \bar{\omega}_{\dot{\alpha}}\partial\bar{\lambda}^{\dot{\alpha}} + T_C - b\partial c. \end{aligned} \quad (5.7)$$

It should be noted that the coefficient of the $\log\pi^+$ term is modified such that the total central charge vanishes. From this form of the energy-momentum tensor, the lower-dimensional

superstring seems to be coupled with the CY sector without twisting. However, if we modify the similarity transformation as $Y \rightarrow Y + Y_C$ with

$$Y_C = - \oint \frac{dz}{2\pi i} J_C \log \left(\frac{2\tilde{\lambda}^2}{\sqrt{\pi^+}} \right) \quad (5.8a)$$

or $\bar{Y} \rightarrow \bar{Y} + \bar{Y}_C$ with

$$\bar{Y}_C = \oint \frac{dz}{2\pi i} J_C \log \left(\frac{2\tilde{\lambda}^2}{\sqrt{\pi^+}} \right), \quad (5.8b)$$

the total BRST charge becomes

$$Q^{tot} = Q + \oint \frac{dz}{2\pi i} G^+, \quad (5.9a)$$

with the BRST charge Q (3.58), for the branch $\bar{\lambda}^\alpha = 0$ and

$$Q^{tot} = Q + \oint \frac{dz}{2\pi i} G^-, \quad (5.9b)$$

with the BRST charge Q (3.67), for the branch $\lambda^\alpha = 0$. The energy-momentum tensor in the CY sector is again twisted as $T_C + \frac{1}{2}\partial J_C$ ($T_C - \frac{1}{2}\partial J_C$) for the branch $\bar{\lambda}^\alpha = 0$ ($\lambda^\alpha = 0$). The cohomology of these BRST charges (5.9a) and (5.9b) are also essentially equivalent to that of the topological charge (5.5).*)

§6. Discussion

In this paper we have investigated the lower-dimensional, $d = 4$ and $d = 6$, superstrings using the DS formalism and shown that they are equivalent to the lower-dimensional PS superstrings. The unexpected off-shell nature of the physical spectrum can be interpreted as a manifestation of noncriticality in the PS formalism.

Because the lower-dimensional PS superstring has no Lorentz anomaly, the symmetry is also realized in the DS formalism, at least in the physical Hilbert space. However, a direct study of Lorentz anomalies in the DS formalism is still worth carrying out, as we have taken the semi-light cone gauge to quantize it. In fact, we have found that the physical Lorentz generator M^{i-} , which is non-trivial in the semi-light cone gauge,^{(17), (18)} does not commute with the BRST charge in lower dimensions. This result will be reported in a forthcoming paper.⁽¹⁹⁾

*) The constraint algebra of the six-dimensional case can also be extended if we impose the $N = 4$ superconformal symmetry on the CY sector. However, it seems there is no similarity transformation to relate it to the PS formalism coupled with the CY sector.

It seems peculiar that the off-shell vector multiplet is included in the physical spectrum, since the DS superstring is classically equivalent with the GS superstring. However, this does not result in a fatal contradiction, since the lower-dimensional GS superstring is not well-defined, due to the global Lorentz anomaly. In addition, the massless spectrum of the GS superstring in the light-cone gauge involves only half of the vector multiplet, which has only positive or negative helicities; there is yet no principle that allows us to use two multiplets with both positive and negative helicities simultaneously. In any case, it is still mysterious why the off-shell states appear as physical states, since in the initial stage we have the Virasoro constraint, which comes from the world-sheet reparametrization invariance. This should be clarified in a future investigation.

It would be interesting to consider how to couple the lower-dimensional DS superstring with the compactified space degrees of freedom. We have proposed two ways to combine the two sectors, although they are not completely successful. It would also be interesting to study the relation to the hybrid formalism,¹³⁾ and this may provide some information about how to make the lower-dimensional superstring consistent. It may also be possible for lower-dimensional superstrings to provide a framework to construct some *off-shell* superstring which is different from the strings in the conventional string theory.^{14)–16)} It would be interesting to explore such a possibility.

Acknowledgements

The authors would like to thank Yoichi Kazama for useful discussions. The work of HK is supported in part by a Grant-in-Aid for Scientific Research (No. 13135213) and a Grant-in-Aid for the 21st Century COE “Center for Diversity and Universality in Physics,” while the work of SM is supported by a Grant-in-Aid for Scientific Research (No. 16540273) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

Appendix A

—— Notation and Conventions ——

In this appendix we present our notation and conventions. We use the metric $\eta^{\mu\nu} = \text{diag}(+1, -1, \dots, -1)$. The light-cone coordinate $x^\mu = (x^\pm, x^i)$ ($i = 1, \dots, d-2$) is defined by $x^\pm = x^0 \pm x^{d-1}$. The gamma matrices satisfy

$$\{\Gamma^\mu, \Gamma^\nu\} = 2\eta^{\mu\nu}. \quad (\text{A}\cdot 1)$$

We use the two-(four-)component spinor notation for $d = 4$ ($d = 6$), employing the following explicit representations of the gamma matrices.

A.1. $d = 4$

We use the two-component notation for the gamma matrices

$$\Gamma_\mu = \begin{pmatrix} 0 & (\sigma_\mu)_{\alpha\dot{\beta}} \\ (\bar{\sigma}_\mu)^{\dot{\alpha}\beta} & 0 \end{pmatrix}, \quad \Gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (\text{A.2})$$

where $\alpha, \beta, \dot{\alpha}, \dot{\beta} = 1, 2$ and

$$(\sigma_\mu)_{\alpha\dot{\beta}} = (1, \boldsymbol{\sigma}), \quad (\bar{\sigma}_\mu)^{\dot{\alpha}\beta} = (1, -\boldsymbol{\sigma}). \quad (\text{A.3})$$

Here, $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ represents the Pauli matrices. These are related as $(\bar{\sigma}^\mu)^{\dot{\alpha}\alpha} = \epsilon^{\alpha\beta}\epsilon^{\dot{\alpha}\dot{\beta}}(\sigma^\mu)_{\beta\dot{\beta}}$, where the two-dimensional antisymmetric two-spinor is defined by $\epsilon^{\alpha\beta} = \epsilon_{\alpha\beta} = \epsilon^{\dot{\alpha}\dot{\beta}} = \epsilon_{\dot{\alpha}\dot{\beta}} = i\sigma_2$. The useful Fierz identities are given by

$$(\sigma^\mu)_{\alpha\dot{\beta}}(\bar{\sigma}_\mu)^{\dot{\gamma}\delta} = 2\delta_\alpha^\delta\delta_{\dot{\beta}}^{\dot{\gamma}}, \quad (\sigma^\mu)_{\alpha\dot{\beta}}(\sigma_\mu)_{\gamma\dot{\delta}} = 2\epsilon_{\alpha\gamma}\epsilon_{\dot{\beta}\dot{\delta}}. \quad (\text{A.4})$$

The indices of the two-component spinors are raised and lowered as

$$\theta^\alpha = \epsilon^{\alpha\beta}\theta_\beta, \quad \theta_\alpha = \theta^\beta\epsilon_{\beta\alpha}, \quad (\text{A.5a})$$

$$\bar{\theta}^{\dot{\alpha}} = \epsilon^{\dot{\alpha}\dot{\beta}}\bar{\theta}_{\dot{\beta}}, \quad \bar{\theta}_{\dot{\alpha}} = \bar{\theta}^{\dot{\beta}}\epsilon_{\dot{\beta}\dot{\alpha}}. \quad (\text{A.5b})$$

In this notation, a Majorana spinor is represented by the pair of complex conjugate Weyl spinors $(\theta^\alpha)^* = \bar{\theta}^{\dot{\alpha}}$.

A.2. $d = 6$

The four-component notation in six dimensions is that employing the representation of 8×8 gamma matrices

$$\Gamma_\mu = \begin{pmatrix} 0 & \bar{\gamma}_\mu \\ \gamma_\mu & 0 \end{pmatrix}, \quad \Gamma_7 = \Gamma_0\Gamma_1\Gamma_2\Gamma_3\Gamma_4\Gamma_5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (\text{A.6})$$

where

$$\gamma_\mu = (\gamma_0, \gamma_i, \gamma_5), \quad \bar{\gamma}_\mu = (\gamma_0, -\gamma_i, -\gamma_5) \quad (i = 1, \dots, 4) \quad (\text{A.7})$$

are 4×4 matrices whose forms we choose as follows:

$$\begin{aligned} \gamma_0 &= 1 \otimes 1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & \gamma_1 &= \sigma_2 \otimes \sigma_1 = \begin{pmatrix} 0 & -i\sigma_1 \\ i\sigma_1 & 0 \end{pmatrix}, \\ \gamma_2 &= \sigma_2 \otimes \sigma_2 = \begin{pmatrix} 0 & -i\sigma_2 \\ i\sigma_2 & 0 \end{pmatrix}, & \gamma_3 &= \sigma_2 \otimes \sigma_3 = \begin{pmatrix} 0 & -i\sigma_3 \\ i\sigma_3 & 0 \end{pmatrix}, \\ \gamma_4 &= \sigma_1 \otimes 1_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, & \gamma_5 &= \sigma_3 \otimes 1_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \end{aligned} \quad (\text{A.8})$$

The charge conjugation matrix satisfying

$$\Gamma^{\mu T} = -\mathcal{C}\Gamma^\mu\mathcal{C}^{-1}, \quad (\text{A}\cdot 9)$$

$$\mathcal{C}^T = \mathcal{C} \quad (\text{A}\cdot 10)$$

is given in this representation by

$$\begin{aligned} \mathcal{C} &= \Gamma^2 \Gamma^4 \Gamma^5 = i\sigma_2 \otimes 1_2 \otimes i\sigma_2, \\ &= \begin{pmatrix} 0 & C \\ C^T & 0 \end{pmatrix}, \end{aligned} \quad (\text{A}\cdot 11)$$

with

$$\begin{aligned} C &= 1_2 \otimes i\sigma_2, \\ &= \begin{pmatrix} i\sigma_2 & 0 \\ 0 & i\sigma_2 \end{pmatrix} = -C^T. \end{aligned} \quad (\text{A}\cdot 12)$$

From (A.9), γ^μ and $\bar{\gamma}^\mu$ satisfy

$$(\gamma^\mu)^T = C\gamma^\mu C^{-1}, \quad (\bar{\gamma}^\mu)^T = C\bar{\gamma}^\mu C^{-1}. \quad (\text{A}\cdot 13)$$

Using this charge conjugation matrix, it is natural to consider the combinations

$$\mathcal{C}\Gamma_\mu = \begin{pmatrix} (C\gamma_\mu)_{\alpha\beta} & 0 \\ 0 & -(C\bar{\gamma}_\mu)^{\dot{\alpha}\dot{\beta}} \end{pmatrix}, \quad (\text{A}\cdot 14)$$

$$\Gamma_\mu \mathcal{C}^{-1} = \begin{pmatrix} (\bar{\gamma}_\mu C^{-1})^{\alpha\beta} & 0 \\ 0 & -(\gamma_\mu C^{-1})_{\dot{\alpha}\dot{\beta}} \end{pmatrix}. \quad (\text{A}\cdot 15)$$

All of these 4×4 matrices are anti-symmetric, *i.e.*

$$(C\gamma^\mu)_{\alpha\beta} = -(C\gamma^\mu)_{\beta\alpha}, \quad (C\bar{\gamma}^\mu)^{\dot{\alpha}\dot{\beta}} = -(C\bar{\gamma}^\mu)^{\dot{\beta}\dot{\alpha}}, \quad (\text{A}\cdot 16a)$$

$$(\bar{\gamma}^\mu C^{-1})^{\alpha\beta} = -(\bar{\gamma}^\mu C^{-1})^{\beta\alpha}, \quad (\gamma^\mu C^{-1})_{\dot{\alpha}\dot{\beta}} = -(\gamma^\mu C^{-1})_{\dot{\beta}\dot{\alpha}}, \quad (\text{A}\cdot 16b)$$

and are related as

$$\frac{1}{2}\epsilon^{\alpha\beta\gamma\delta}(C\gamma^\mu)_{\gamma\delta} = -(\bar{\gamma}^\mu C^{-1})^{\alpha\beta}, \quad \frac{1}{2}\epsilon_{\alpha\beta\gamma\delta}(\bar{\gamma}^\mu C^{-1})^{\gamma\delta} = -(C\gamma^\mu)_{\alpha\beta}. \quad (\text{A}\cdot 17)$$

An important Fierz identity is given by

$$\delta_\gamma^{[\alpha}\delta_\delta^{\beta]} = \frac{1}{2}(\delta_\gamma^\alpha\delta_\delta^\beta - \delta_\gamma^\beta\delta_\delta^\alpha) = -\frac{1}{4}(C\gamma^\mu)_{\gamma\delta}(\bar{\gamma}_\mu C^{-1})^{\alpha\beta}, \quad (\text{A}\cdot 18)$$

which can be rewritten using Eq. (A·17) as

$$(C\gamma_\mu)_{\alpha\beta}(C\gamma^\mu)_{\gamma\delta} = 2\epsilon_{\alpha\beta\gamma\delta}, \quad (\bar{\gamma}_\mu C^{-1})^{\alpha\beta}(\bar{\gamma}^\mu C^{-1})^{\gamma\delta} = 2\epsilon^{\alpha\beta\gamma\delta}, \quad (\text{A}\cdot 19)$$

or equivalently

$$(C\gamma_\mu)_{\alpha(\beta}(C\gamma^\mu)_{\gamma)\delta} = 0, \quad (\bar{\gamma}_\mu C^{-1})^{\alpha(\beta}(\bar{\gamma}^\mu C^{-1})^{\gamma)\delta} = 0. \quad (\text{A}\cdot 20)$$

The $SU(2)$ -Majorana-Weyl (MW) spinor θ_{+I} ($I = 1, 2$) satisfies the conditions

$$\Gamma_7 \theta_{+I} = \theta_{+I}, \quad (\text{A}\cdot 21a)$$

$$\bar{\theta}_+^I = \epsilon^{IJ} \theta_{+J}^T \mathcal{C}, \quad (\text{A}\cdot 21b)$$

where ϵ^{IJ} is the $SU(2)$ anti-symmetric tensor defined by $\epsilon^{12} = \epsilon_{12} = 1$, which is used to raise or lower the index I using the same rule as in the case of the four-dimensional spinor (A·5). This $SU(2)$ MW spinor can be written in the four-component notation as

$$\theta_{+I} = \begin{pmatrix} \theta_I^\alpha \\ 0 \end{pmatrix}, \quad (\theta_I^\alpha)^\dagger = \epsilon^{IJ} (\theta_J^T C)^\alpha. \quad (\text{A}\cdot 22)$$

We can further decompose it with the light-cone decomposition

$$\theta_I^\alpha = \begin{pmatrix} \theta_I^a \\ \theta_I^{\dot{a}} \end{pmatrix}, \quad (a, \dot{a} = 1, 2) \quad (\text{A}\cdot 23)$$

where a and $\dot{a}$ are the spinor indices of the transverse rotation $SO(4) \sim SU(2) \times SU(2)$. The $SU(2)$ MW condition is now simply given by

$$(\theta_I^a)^* = \epsilon^{IJ} \theta_J^b \epsilon_{ba} \equiv \theta_a^I, \quad (\text{A}\cdot 24a)$$

$$(\theta_I^{\dot{a}})^* = \epsilon^{IJ} \theta_J^{\dot{b}} \epsilon_{\dot{b}\dot{a}} \equiv \theta_{\dot{a}}^I. \quad (\text{A}\cdot 24b)$$

From the representation (A·8), the transverse components of the gamma matrices can also be decomposed using the 2×2 matrices $\tilde{\gamma}_i$ defined by

$$\tilde{\gamma}_i = i\sigma_i \quad (i = 1, 2, 3), \quad \tilde{\gamma}_4 = 1_2. \quad (\text{A}\cdot 25)$$

It is also useful to define $\tilde{\tilde{\gamma}}_i$ as

$$\tilde{\tilde{\gamma}}_i = -i\sigma_i \quad (i = 1, 2, 3), \quad \tilde{\tilde{\gamma}}_4 = 1_2. \quad (\text{A}\cdot 26)$$

These two 2×2 matrices are related as

$$(\epsilon \tilde{\gamma}_i)_{\dot{a}b} = -(\epsilon \tilde{\tilde{\gamma}}_i)_{b\dot{a}}, \quad (\text{A}\cdot 27)$$

where $\epsilon = i\sigma_2$.

The two quantities $\mathbb{I}$ and $\bar{\mathbb{I}}$ constitute a representation of the quaternion and its conjugate, and they satisfy the relation

$$\mathbb{I} \bar{\mathbb{I}} = \bar{\mathbb{I}} \mathbb{I} = \Pi^i \Pi^i. \quad (\text{A}\cdot 28)$$

References

- 1) N. Berkovits, J. High Energy Phys. **04** (2000), 018; hep-th/0001035.
- 2) P. A. Grassi and N. Wyllard, J. High Energy Phys. **12** (2005), 007; hep-th/0509140.
- 3) N. Wyllard, J. High Energy Phys. **11** (2005), 009; hep-th/0509165.
- 4) Y. Aisaka and Y. Kazama, J. High Energy Phys. **05** (2005), 046; hep-th/0502208.
- 5) J. Wess and J. Bagger, “*Supersymmetry and Supergravity*” (Princeton University Press, 1992).
- 6) P. S. Howe, G. Sierra and P. K. Townsend, Nucl. Phys. B **221** (1983), 331.
- 7) M. B. Green and J. H. Schwarz, Phys. Lett. B **136** (1984), 367.
- 8) Y. Aisaka and Y. Kazama, J. High Energy Phys. **02** (2003), 017; hep-th/0212316.
- 9) N. Berkovits and C. Vafa, Mod. Phys. Lett. A **9** (1994), 653; hep-th/9310170.
- 10) N. Ohta and J. L. Petersen, Phys. Lett. B **325** (1994), 67; hep-th/9312187.
- 11) O. Chandia, J. High Energy Phys. **10** (2005), 105; hep-th/0509185.
- 12) N. Berkovits, J. High Energy Phys. **10** (2005), 089; hep-th/0509120.
- 13) N. Berkovits, Nucl. Phys. B **431** (1994), 258; hep-th/9404162.
- 14) A. Giveon, D. Kutasov and O. Pelc, J. High Energy Phys. **10** (1999), 035; hep-th/9907178.
- 15) S. Mizoguchi, J. High Energy Phys. **04** (2000), 014; hep-th/0003053.
- 16) I. Adam, P. A. Grassi, L. Mazzucato, Y. Oz and S. Yankielowicz, “*Non-Critical Pure Spinor Superstrings*”, hep-th/0605118.
- 17) N. Berkovits and D. Z. Marchioro, J. High Energy Phys. **01** (2005), 018; hep-th/0412198.
- 18) A. Gaora and J. A. Garcia, J. High Energy Phys. **09** (2005), 083; hep-th/0507076.
- 19) H. Kunitomo and S. Mizoguchi, in preparation.